

# **UVRAG Antibody (C-Terminus)**

Rabbit Polyclonal Antibody Catalog # ALS16426

# **Specification**

## **UVRAG Antibody (C-Terminus) - Product Information**

Application IF

Primary Accession <u>Q9P2Y5</u>

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Calculated MW 78kDa KDa

### **UVRAG Antibody (C-Terminus) - Additional Information**

### **Gene ID 7405**

#### **Other Names**

UV radiation resistance-associated gene protein, p63, UVRAG

# Target/Specificity

UVRAG antibody is human and mouse reactive.

### **Reconstitution & Storage**

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

#### **Precautions**

UVRAG Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

# **UVRAG Antibody (C-Terminus) - Protein Information**

### Name UVRAG

#### **Function**

Versatile protein that is involved in regulation of different cellular pathways implicated in membrane trafficking. Involved in regulation of the COPI-dependent retrograde transport from Golgi and the endoplasmic reticulum by associating with the NRZ complex; the function is dependent on its binding to phosphatidylinositol 3- phosphate (PtdIns(3)P) (PubMed:<a href="http://www.uniprot.org/citations/16799551" target="\_blank">16799551</a>, PubMed:<a href="http://www.uniprot.org/citations/18552835" target="\_blank">18552835</a>, PubMed:<a href="http://www.uniprot.org/citations/20643123" target="\_blank">20643123</a>, PubMed:<a href="http://www.uniprot.org/citations/24056303" target="\_blank">24056303</a>, PubMed:<a href="http://www.uniprot.org/citations/28306502" target="\_blank">28306502</a>). During autophagy acts as a regulatory subunit of the alternative Pl3K complex II (Pl3KC3-C2) that mediates formation of phosphatidylinositol 3-phosphate and is believed to be involved in maturation of autophagosomes and endocytosis. Activates lipid kinase activity of PlK3C3 (PubMed:<a href="http://www.uniprot.org/citations/16799551" target="\_blank">16799551</a>/a>,



PubMed:<a href="http://www.uniprot.org/citations/20643123" target=" blank">20643123</a>, PubMed: <a href="http://www.uniprot.org/citations/24056303" target="blank">24056303</a>,  $PubMed: <a href="http://www.uniprot.org/citations/28306502" target="\_blank">28306502 </a>).$ Involved in the regulation of degradative endocytic trafficking and cytokinesis, and in regulation of ATG9A transport from the Golgi to the autophagosome; the functions seems to implicate its association with PI3KC3-C2 (PubMed: <a href="http://www.uniprot.org/citations/16799551" target=" blank">16799551</a>, PubMed:<a href="http://www.uniprot.org/citations/20643123" target="blank">20643123</a>, PubMed:<a href="http://www.uniprot.org/citations/24056303" target="blank">24056303</a>). Involved in maturation of autophagosomes and degradative endocytic trafficking independently of BECN1 but depending on its association with a class C Vps complex (possibly the HOPS complex); the association is also proposed to promote autophagosome recruitment and activation of Rab7 and endosome-endosome fusion events (PubMed:<a href="http://www.uniprot.org/citations/18552835" target=" blank">18552835</a>, PubMed:<a href="http://www.uniprot.org/citations/28306502" target=" blank">28306502</a>). Enhances class C Vps complex (possibly HOPS complex) association with a SNARE complex and promotes fusogenic SNARE complex formation during late endocytic membrane fusion (PubMed:<a href="http://www.uniprot.org/citations/24550300" target=" blank">24550300</a>). In case of negative- strand RNA virus infection is required for efficient virus entry, promotes endocytic transport of virions and is implicated in a VAMP8- specific fusogenic SNARE complex assembly (PubMed:<a href="http://www.uniprot.org/citations/24550300" target=" blank">24550300</a>).

#### **Cellular Location**

Late endosome. Lysosome. Cytoplasmic vesicle, autophagosome. Early endosome. Endoplasmic reticulum. Midbody. Chromosome, centromere. Note=Colocalizes with RAB9-positive compartments involved in retrograde transport from late endosomes to trans-Golgi network. Colocalization with early endosomes is only partial (PubMed:24056303). Recruited to autophagosome following interaction with RUBCNL/PACER (PubMed:28306502)

## **Tissue Location**

Highly expressed in brain, lung, kidney and liver.


# **UVRAG Antibody (C-Terminus) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **UVRAG Antibody (C-Terminus) - Images**





Immunofluorescence of UVRAG in human lung tissue with UVRAG antibody at 20 ug/ml.

# **UVRAG Antibody (C-Terminus) - References**

Perelman B.,et al.Genomics 41:397-405(1997). lida A.,et al.Hum. Genet. 106:277-287(2000). Ota T.,et al.Nat. Genet. 36:40-45(2004). Taylor T.D.,et al.Nature 440:497-500(2006). ltakura E.,et al.Mol. Biol. Cell 19:5360-5372(2008).