10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BCL3 / BCL-3 Antibody (C-Terminus)

Rabbit Polyclonal Antibody

Catalog # ALS16743

Specification

BCL3 / BCL-3 Antibody (C-Terminus) - Product Information

Application	**IHC, WB**
Primary Accession	P20749
Other Accession	602
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**IgG**
Calculated MW	**47584**

BCL3 / BCL-3 Antibody (C-Terminus) - Additional Information

Gene ID 602

Other Names
BCL3, B-cell CLL/lymphoma 3, B-cell leukemia/lymphoma 3, B-cell lymphoma 3 protein, BCL-3, BCL4, D19S37, Proto-oncogene BCL3

Target/Specificity
Human BCL3 / BCL-3

Reconstitution & Storage
PBS, 0.02% sodium azide. Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions
BCL3 / BCL-3 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

BCL3 / BCL-3 Antibody (C-Terminus) - Protein Information

Name BCL3

Synonyms BCL4, D19S37

Function
Contributes to the regulation of transcriptional activation of NF-kappa-B target genes. In the cytoplasm, inhibits the nuclear translocation of the NF-kappa-B p50 subunit. In the nucleus, acts as transcriptional activator that promotes transcription of NF-kappa-B target genes. Contributes to the regulation of cell proliferation (By similarity).

Cellular Location
Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Note=Ubiquitination via 'Lys-63'- linked ubiquitin chains is required for nuclear accumulation


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BCL3 / BCL-3 Antibody (C-Terminus) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

BCL3 / BCL-3 Antibody (C-Terminus) - Images



Anti-BCL3 / BCL-3 antibody IHC staining of human tonsil.



Western blot analysis of BCL3 in 293 cell lysate with BCL3 antibody at 1 ug/ml.

BCL3 / BCL-3 Antibody (C-Terminus) - Background

Contributes to the regulation of transcriptional activation of NF-kappa-B target genes. In the cytoplasm, inhibits the nuclear translocation of the NF-kappa-B p50 subunit. In the nucleus, acts as transcriptional activator that promotes transcription of NF-kappa-B target genes. Contributes to the regulation of cell proliferation (By similarity).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BCL3 / BCL-3 Antibody (C-Terminus) - References

Ohno H.,et al.Cell 60:991-997(1990).
McKeithan T.W.,et al.Genomics 24:120-126(1994).
Bours V.,et al.Cell 72:729-739(1993).
Heissmeyer V.,et al.EMBO J. 18:4766-4778(1999).
Dechend R.,et al.Oncogene 18:3316-3323(1999).