

BATF3 Antibody (N-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS17112**Specification**

BATF3 Antibody (N-Terminus) - Product Information

Application	IHC
Primary Accession	Q9NR55
Other Accession	55509
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	14468

BATF3 Antibody (N-Terminus) - Additional Information**Gene ID** 55509**Other Names**

BATF3, JUNDM1, JDP1, Jun dimerization protein 1, B-ATF-3, SNFT

Target/Specificity

BATF3 antibody is human, mouse and rat reactive. BATF3 antibody is predicted to have no cross-reactivity to other BATF family members.

Reconstitution & Storage

PBS, 0.02% sodium azide. Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions

BATF3 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

BATF3 Antibody (N-Terminus) - Protein Information**Name** BATF3**Synonyms** SNFT**Function**

AP-1 family transcription factor that controls the differentiation of CD8(+) thymic conventional dendritic cells in the immune system. Required for development of CD8-alpha(+) classical dendritic cells (cDCs) and related CD103(+) dendritic cells that cross-present antigens to CD8 T-cells and produce interleukin-12 (IL12) in response to pathogens (By similarity). Acts via the formation of a heterodimer with JUN family proteins that recognizes and binds DNA sequence 5'-TGA[CG]TCA-3' and regulates expression of target genes.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00978, ECO:0000269|PubMed:12087103}

BATF3 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BATF3 Antibody (N-Terminus) - Images



Human Skin: Formalin-Fixed, Paraffin-Embedded (FFPE)

BATF3 Antibody (N-Terminus) - Background

AP-1 family transcription factor that controls the differentiation of CD8(+) thymic conventional dendritic cells in the immune system. Required for development of CD8-alpha(+) classical dendritic cells (cDCs) and related CD103(+) dendritic cells that cross-present antigens to CD8 T-cells and produce interleukin-12 (IL12) in response to pathogens (By similarity). Acts via the formation of a heterodimer with JUN family proteins that recognizes and binds DNA sequence 5'-TGA[CG]TCA-3' and regulates expression of target genes.

BATF3 Antibody (N-Terminus) - References

Iacobelli M., et al. J. Immunol. 165:860-868(2000).
Gregory S.G., et al. Nature 441:315-321(2006).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Bower K.E., et al. J. Biol. Chem. 277:34967-34977(2002).
Bower K.E., et al. Oncogene 23:8805-8814(2004).