

Anti-CST3 / Cystatin C Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17282**Specification**

Anti-CST3 / Cystatin C Antibody - Product Information

Application	IHC-P
Primary Accession	P01034
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	15799

Anti-CST3 / Cystatin C Antibody - Additional Information**Gene ID** 1471**Alias Symbol** CST3**Other Names**

CST3, ARMD11, BA218C14.4 (cystatin C), Cystatin-3, Cystatin 3, Cystatin C, Cystatin-C, Gamma-trace, Post-gamma-globulin

Target/Specificity

Human CST3 / Cystatin C

Reconstitution & Storage

PBS, pH 7.4, 0.03% Proclin 300, 50% glycerol. Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions

Anti-CST3 / Cystatin C Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-CST3 / Cystatin C Antibody - Protein Information**Name** CST3**Function**

As an inhibitor of cysteine proteinases, this protein is thought to serve an important physiological role as a local regulator of this enzyme activity.

Cellular Location

Secreted.

Tissue LocationExpressed in submandibular and sublingual saliva but not in parotid saliva (at protein level).
Expressed in various body fluids, such as the cerebrospinal fluid and plasma. Expressed in highest


levels in the epididymis, vas deferens, brain, thymus, and ovary and the lowest in the submandibular gland

Anti-CST3 / Cystatin C Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CST3 / Cystatin C Antibody - Images



Human Kidney: Formalin-Fixed, Paraffin-Embedded (FFPE)