

Anti-IFNAR2 Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17356**Specification**

Anti-IFNAR2 Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P48551
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	57759

Anti-IFNAR2 Antibody - Additional Information**Gene ID** 3455**Alias Symbol** IFNAR2**Other Names**

IFNAR2, IFN-alpha/beta receptor 2, IFN-alpha-REC, IFN-alpha binding protein, IFN-R, IFN-R-2, IFNARB, Type I interferon receptor 2, IFNABR

Target/Specificity

Human IFNAR2

Reconstitution & Storage

PBS, pH 7.3, 0.02% sodium azide, 50% glycerol. Long term: -80°C; Short term: -20°C. Avoid freeze-thaw cycles.

Precautions

Anti-IFNAR2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-IFNAR2 Antibody - Protein Information**Name** IFNAR2**Synonyms** IFNABR, IFNARB**Function**

Together with IFNAR1, forms the heterodimeric receptor for type I interferons (including interferons alpha, beta, epsilon, omega and kappa) (PubMed:8181059, PubMed:7665574, PubMed:7759950, PubMed:8798579, PubMed:8969169, PubMed:8969169)

href="http://www.uniprot.org/citations/10049744" target="_blank">10049744, PubMed:10556041, PubMed:21854986, PubMed:26424569, PubMed:28165510, PubMed:32972995). Type I interferon binding activates the JAK-STAT signaling cascade, resulting in transcriptional activation or repression of interferon-regulated genes that encode the effectors of the interferon response (PubMed:8181059, PubMed:7665574, PubMed:7759950, PubMed:8798579, PubMed:8969169, PubMed:10049744, PubMed:17517919, PubMed:21854986, PubMed:26424569, PubMed:28165510, PubMed:32972995). Mechanistically, type I interferon-binding brings the IFNAR1 and IFNAR2 subunits into close proximity with one another, driving their associated Janus kinases (JAKs) (TYK2 bound to IFNAR1 and JAK1 bound to IFNAR2) to cross-phosphorylate one another (PubMed:10556041, PubMed:11682488, PubMed:12105218, PubMed:21854986, PubMed:32972995). The activated kinases phosphorylate specific tyrosine residues on the intracellular domains of IFNAR1 and IFNAR2, forming docking sites for the STAT transcription factors (STAT1, STAT2 and STAT3) (PubMed:11682488, PubMed:12105218, PubMed:21854986, PubMed:32972995). STAT proteins are then phosphorylated by the JAKs, promoting their translocation into the nucleus to regulate expression of interferon-regulated genes (PubMed:9121453, PubMed:12105218, PubMed:28165510).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 3]: Secreted

Tissue Location

Isoform 3 is detected in the urine (at protein level) (PubMed:8181059, PubMed:7759950).
Expressed in blood cells Expressed in lymphoblastoid and fibrosarcoma cell lines

Anti-IFNAR2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IFNAR2 Antibody - Images