

Anti-COMP / THBS5 Antibody

Rabbit Anti Human Polyclonal Antibody Catalog # ALS17366

Specification

Anti-COMP / THBS5 Antibody - Product Information

Application WB, IHC-P Primary Accession P49747

Predicted Human, Mouse, Rat

Host Rabbit Clonality Polyclonal

Isotype IgG
Calculated MW 82860

Anti-COMP / THBS5 Antibody - Additional Information

Gene ID 1311

Alias Symbol COMP

Other Names

COMP, MED, PSACH, Thrombospondin-5, TSP5, THBS5, EDM1, EPD1

Target/Specificity

Human COMP / THBS5.

Reconstitution & Storage

PBS, pH 7.3, 0.02% sodium azide, 50% glycerol. Long term: -80°C; Short term: -20°C. Avoid freeze-thaw cycles.

Precautions

Anti-COMP / THBS5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-COMP / THBS5 Antibody - Protein Information

Name COMP (HGNC:2227)

Function

Plays a role in the structural integrity of cartilage via its interaction with other extracellular matrix proteins such as the collagens and fibronectin. Can mediate the interaction of chondrocytes with the cartilage extracellular matrix through interaction with cell surface integrin receptors (PubMed:16542502, PubMed:16051604). Could play a role in the pathogenesis of osteoarthritis (PubMed:16542502). Potent

href="http://www.uniprot.org/citations/16542502" target="_blank">16542502). Potent suppressor of apoptosis in both primary chondrocytes and transformed cells. Suppresses apoptosis by blocking the activation of caspase-3 and by inducing the IAP family of survival proteins (BIRC3, BIRC2, BIRC5 and XIAP) (PubMed:<a href="http://www.uniprot.org/citations/17993464"



target="_blank">17993464). Essential for maintaining a vascular smooth muscle cells (VSMCs) contractile/differentiated phenotype under physiological and pathological stimuli. Maintains this phenotype of VSMCs by interacting with ITGA7 (By similarity).

Cellular Location

Secreted, extracellular space, extracellular matrix

Tissue Location

Abundantly expressed in the chondrocyte extracellular matrix, and is also found in bone, tendon, ligament and synovium and blood vessels. Increased amounts are produced during late stages of osteoarthritis in the area adjacent to the main defect

Anti-COMP / THBS5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-COMP / THBS5 Antibody - Images