

Anti-KDM1A / LSD1 Antibody (aa97-328)
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17394**Specification****Anti-KDM1A / LSD1 Antibody (aa97-328) - Product Information**

Application	WB, IHC-P
Primary Accession	O60341
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	92903

Anti-KDM1A / LSD1 Antibody (aa97-328) - Additional Information**Gene ID** 23028

Alias Symbol	KDM1A
Other Names	
KDM1A, BHC110, KDM1, LSD1, AOF2, KIAA0601	

Target/Specificity
Human KDM1A / LSD1**Reconstitution & Storage**

PBS, 0.05% sodium azide. Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Precautions

Anti-KDM1A / LSD1 Antibody (aa97-328) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-KDM1A / LSD1 Antibody (aa97-328) - Protein Information**Name** KDM1A ([HGNC:29079](#))**Function**

Histone demethylase that can demethylate both 'Lys-4' (H3K4me) and 'Lys-9' (H3K9me) of histone H3, thereby acting as a coactivator or a corepressor, depending on the context (PubMed: [15620353](http://www.uniprot.org/citations/15620353), PubMed: [15811342](http://www.uniprot.org/citations/15811342), PubMed: [16140033](http://www.uniprot.org/citations/16140033), PubMed: [16079794](http://www.uniprot.org/citations/16079794), PubMed: [16079795](http://www.uniprot.org/citations/16079795), PubMed: [16223729](http://www.uniprot.org/citations/16223729)). Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed (PubMed: [15620353](http://www.uniprot.org/citations/15620353), PubMed: [15811342](http://www.uniprot.org/citations/15811342)

target="_blank">15811342, PubMed:16079794, PubMed:21300290). Acts as a corepressor by mediating demethylation of H3K4me, a specific tag for epigenetic transcriptional activation. Demethylates both mono- (H3K4me1) and di-methylated (H3K4me2) H3K4me (PubMed:15620353, PubMed:20389281, PubMed:21300290, PubMed:23721412). May play a role in the repression of neuronal genes. Alone, it is unable to demethylate H3K4me on nucleosomes and requires the presence of RCOR1/CoREST to achieve such activity (PubMed:16140033, PubMed:16079794, PubMed:16885027, PubMed:21300290, PubMed:23721412). Also acts as a coactivator of androgen receptor (AR)-dependent transcription, by being recruited to AR target genes and mediating demethylation of H3K9me, a specific tag for epigenetic transcriptional repression. The presence of PRKCB in AR-containing complexes, which mediates phosphorylation of 'Thr-6' of histone H3 (H3T6ph), a specific tag that prevents demethylation H3K4me, prevents H3K4me demethylase activity of KDM1A (PubMed:16079795). Demethylates di-methylated 'Lys-370' of p53/TP53 which prevents interaction of p53/TP53 with TP53BP1 and represses p53/TP53-mediated transcriptional activation. Demethylates and stabilizes the DNA methylase DNMT1 (PubMed:29691401). Demethylates methylated 'Lys-42' and methylated 'Lys-117' of SOX2 (PubMed:29358331). Required for gastrulation during embryogenesis. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. Effector of SNAI1-mediated transcription repression of E-cadherin/CDH1, CDN7 and KRT8. Required for the maintenance of the silenced state of the SNAI1 target genes E-cadherin/CDH1 and CDN7 (PubMed:20389281).

Cellular Location

Nucleus

Tissue Location

Ubiquitously expressed.

Anti-KDM1A / LSD1 Antibody (aa97-328) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-KDM1A / LSD1 Antibody (aa97-328) - Images