10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-CD135 / FLT3 Antibody (Internal)

Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17567

Specification

Anti-CD135 / FLT3 Antibody (Internal) - Product Information

Application	**IHC-P, E**
Primary Accession	P36888
Predicted	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Calculated MW	**112903**

Anti-CD135 / FLT3 Antibody (Internal) - Additional Information

Gene ID 2322

Alias Symbol	**FLT3**

Other Names
FLT3, CD135, CD135 antigen, Fetal liver kinase 2, Fetal liver kinase-2, FL cytokine receptor, FLK2, FLT-3, Fms-like tyrosine kinase 3, Stem cell tyrosine kinase 1, STK-1, FLK-2, Fms-related tyrosine kinase 3, STK1

Target/Specificity
Human FLT3 / CD135. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage
Immunoaffinity purified

Precautions
Anti-CD135 / FLT3 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-CD135 / FLT3 Antibody (Internal) - Protein Information

Name FLT3

Synonyms CD135, FLK2, STK1

Function
Tyrosine-protein kinase that acts as a cell-surface receptor for the cytokine FLT3LG and regulates differentiation, proliferation and survival of hematopoietic progenitor cells and of dendritic cells. Promotes phosphorylation of SHC1 and AKT1, and activation of the downstream effector MTOR. Promotes activation of RAS signaling and phosphorylation of downstream kinases, including MAPK1/ERK2 and/or MAPK3/ERK1. Promotes phosphorylation of FES, FER, PTPN6/SHP, PTPN11/SHP-2, PLCG1, and STAT5A and/or STAT5B. Activation of wild-type FLT3 causes only marginal activation of STAT5A or STAT5B. Mutations that cause constitutive kinase activity

promote cell proliferation and resistance to apoptosis via the activation of multiple signaling pathways.

Cellular Location
Membrane; Single-pass type I membrane protein. Endoplasmic reticulum lumen. Note=Constitutively activated mutant forms with internal tandem duplications are less efficiently transported to the cell surface and a significant proportion is retained in an immature form in the endoplasmic reticulum lumen. The activated kinase is rapidly targeted for degradation

Tissue Location
Detected in bone marrow, in hematopoietic stem cells, in myeloid progenitor cells and in granulocyte/macrophage progenitor cells (at protein level). Detected in bone marrow, liver, thymus, spleen and lymph node, and at low levels in kidney and pancreas. Highly expressed in T-cell leukemia

Anti-CD135 / FLT3 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-CD135 / FLT3 Antibody (Internal) - Images