

Anti-SLC7A11 / XCT Antibody (N-Terminus)
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17576**Specification**

Anti-SLC7A11 / XCT Antibody (N-Terminus) - Product Information

Application	IHC-P
Primary Accession	Q9UPY5
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55423

Anti-SLC7A11 / XCT Antibody (N-Terminus) - Additional Information**Gene ID** 23657

Alias Symbol	SLC7A11
Other Names	
SLC7A11, CCB1, Cystine/glutamate transporter, XCT	

Target/Specificity

Human SLC7A11. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Immunoaffinity purified

Precautions

Anti-SLC7A11 / XCT Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-SLC7A11 / XCT Antibody (N-Terminus) - Protein Information**Name** SLC7A11 ([HGNC:11059](#))**Function**

Heterodimer with SLC3A2, that functions as an antiporter by mediating the exchange of extracellular anionic L-cystine and intracellular L-glutamate across the cellular plasma membrane (PubMed: [15151999](http://www.uniprot.org/citations/15151999), PubMed: [34880232](http://www.uniprot.org/citations/34880232), PubMed: [35352032](http://www.uniprot.org/citations/35352032), PubMed: [35245456](http://www.uniprot.org/citations/35245456), PubMed: [11417227](http://www.uniprot.org/citations/11417227), PubMed: [14722095](http://www.uniprot.org/citations/14722095), PubMed: [11133847](http://www.uniprot.org/citations/11133847)). Provides L-cystine for the maintenance of the redox balance between extracellular L-cystine and L-cysteine and for the maintenance of the intracellular levels of glutathione that is essential for

cells protection from oxidative stress (By similarity). The transport is sodium-independent, electroneutral with a stoichiometry of 1:1, and is driven by the high intracellular concentration of L-glutamate and the intracellular reduction of L-cystine (PubMed:11417227, PubMed:11133847). In addition, mediates the import of L-kynurenine leading to anti-ferroptotic signaling propagation required to maintain L-cystine and glutathione homeostasis (PubMed:35245456). Moreover, mediates N-acetyl-L-cysteine uptake into the placenta leading to subsequently down-regulation of pathways associated with oxidative stress, inflammation and apoptosis (PubMed:34120018). In vitro can also transport L-aspartate (PubMed:11417227). May participate in astrocyte and meningeal cell proliferation during development and can provide neuroprotection by promoting glutathione synthesis and delivery from non-neuronal cells such as astrocytes and meningeal cells to immature neurons (By similarity). Controls the production of pheomelanin pigment directly (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, microvillus membrane; Multi-pass membrane protein. Note=Localized to the microvillous membrane of the placental syncytiotrophoblast.

Tissue Location

Expressed in term placenta and primary term cytotrophoblast (PubMed:34120018). Expressed mainly in the brain, but also in pancreas (PubMed:11417227).

Anti-SLC7A11 / XCT Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SLC7A11 / XCT Antibody (N-Terminus) - Images