

Anti-GJA1 / CX43 / Connexin 43 Antibody (Internal) Rabbit Anti Human Polyclonal Antibody Catalog # ALS17578

Specification

Anti-GJA1 / CX43 / Connexin 43 Antibody (Internal) - Product Information

Application Primary Accession Predicted Host Clonality Calculated MW IHC-P <u>P17302</u> Human, Chicken Rabbit Polyclonal 43008

Anti-GJA1 / CX43 / Connexin 43 Antibody (Internal) - Additional Information

Gene ID 2697

Alias Symbol

GJA1

Other Names GJA1, Connexin-43, DFNB38, Connexin 43, Gap junction alpha-1 protein, GJAL, HSS, Gap junction protein alpha 1, ODDD, ODD, AVSD3, CX43, HLHS1, ODOD, SDTY3

Target/Specificity Human GJA1 / Connexin 43. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage Immunoaffinity purified

Precautions Anti-GJA1 / CX43 / Connexin 43 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-GJA1 / CX43 / Connexin 43 Antibody (Internal) - Protein Information

Name GJA1

Synonyms GJAL

Function

Gap junction protein that acts as a regulator of bladder capacity. A gap junction consists of a cluster of closely packed pairs of transmembrane channels, the connexons, through which materials of low MW diffuse from one cell to a neighboring cell. May play a critical role in the physiology of hearing by participating in the recycling of potassium to the cochlear endolymph. Negative regulator of bladder functional capacity: acts by enhancing intercellular electrical and chemical transmission, thus sensitizing bladder muscles to cholinergic neural stimuli and causing them to contract (By similarity). May play a role in cell growth inhibition through the regulation of NOV expression and localization. Plays an essential role in gap junction communication in the



ventricles (By similarity).

Cellular Location Cell membrane; Multi-pass membrane protein. Cell junction, gap junction. Endoplasmic reticulum {ECO:0000250|UniProtKB:P23242}. Note=Localizes at the intercalated disk (ICD) in cardiomyocytes and the proper localization at ICD is dependent on TMEM65. {ECO:0000250|UniProtKB:P23242}

Tissue Location Expressed in the heart and fetal cochlea.

Anti-GJA1 / CX43 / Connexin 43 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-GJA1 / CX43 / Connexin 43 Antibody (Internal) - Images