

**Anti-TLR4 Antibody (aa420-435)  
Rabbit Anti Human Polyclonal Antibody  
Catalog # ALS17778**

## Specification

## Anti-TLR4 Antibody (aa420-435) - Product Information

Application	WB, IHC-P, ICC
Primary Accession	<a href="#"><u>000206</u></a>
Predicted	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	95680

## Anti-TLR4 Antibody (aa420-435) - Additional Information

Gene ID 7099

Alias Symbol	TLR4
Other Names	

TLR4, CD284, CD284 antigen, Homolog of Drosophila toll, TLR-4, TOLL, ARMD10, Toll-like receptor 4, HToll

## Target/Specificity

Detects ~75-80kD when tested against partial recombinant mouse TLR4 (extracellular portion plus His-tag).

## **Reconstitution & Storage**

### Affinity purified

## Precautions

**Precautions:** Anti-TLR4 Antibody (aa420-435) is for research use only and not for use in diagnostic or therapeutic procedures.

## Anti-TLR4 Antibody (aa420-435) - Protein Information

Name TLR4

## Function

At the plasma membrane, cooperates with LY96 to mediate the innate immune response to bacterial lipopolysaccharide (LPS) (PubMed:<a href="http://www.uniprot.org/citations/27022195" target="\_blank">27022195</a>, PubMed:<a href="http://www.uniprot.org/citations/17292937" target="\_blank">17292937</a>, PubMed:<a href="http://www.uniprot.org/citations/29038465" target="\_blank">29038465</a>). Also involved in LPS-independent inflammatory responses triggered by free fatty acids, such as palmitate, and Ni(2+) (PubMed:<a href="http://www.uniprot.org/citations/20711192" target="\_blank">20711192</a>). Mechanistically, acts via MYD88, TIRAP and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:<a href="http://www.uniprot.org/citations/9237759" target="\_blank">9237759</a>, PubMed:<a href="http://www.uniprot.org/citations/10835634" target="\_blank">10835634</a>, PubMed:<a href="http://www.uniprot.org/citations/27022195" target="\_blank">27022195</a>, PubMed:<a href="http://www.uniprot.org/citations/21393102" target="\_blank">21393102</a>). Alternatively, CD14-mediated TLR4 internalization via endocytosis is associated with the initiation of a MYD88-independent signaling via the TICAM1-TBK1-IRF3 axis leading to type I interferon production (PubMed:<a href="http://www.uniprot.org/citations/14517278" target="\_blank">14517278</a>). In addition to the secretion of proinflammatory cytokines, initiates the activation of NLRP3 inflammasome and formation of a positive feedback loop between autophagy and NF-kappa-B signaling cascade (PubMed:<a href="http://www.uniprot.org/citations/32894580" target="\_blank">32894580</a>). In complex with TLR6, promotes inflammation in monocytes/macrophages by associating with TLR6 and the receptor CD86 (PubMed:<a href="http://www.uniprot.org/citations/23880187" target="\_blank">23880187</a>). Upon ligand binding, such as oxLDL or amyloid-beta 42, the TLR4:TLR6 complex is internalized and triggers inflammatory response, leading to NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/23880187" target="\_blank">23880187</a>). In myeloid dendritic cells, vesicular stomatitis virus glycoprotein G but not LPS promotes the activation of IRF7, leading to type I IFN production in a CD14-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/23880187" target="\_blank">23880187</a>, PubMed:<a href="http://www.uniprot.org/citations/15265881" target="\_blank">15265881</a>). Required for the migration-promoting effects of ZG16B/PAUF on pancreatic cancer cells.

### Cellular Location

Cell membrane; Single-pass type I membrane protein. Early endosome. Cell projection, ruffle {ECO:0000250|UniProtKB:Q9QUK6}. Note=Upon complex formation with CD36 and TLR6, internalized through dynamin-dependent endocytosis (PubMed:20037584). Colocalizes with RFTN1 at cell membrane and then together with RFTN1 moves to endosomes, upon lipopolysaccharide stimulation. Co-localizes with ZG16B/PAUF at the cell membrane of pancreatic cancer cells (PubMed:36232715)

### Tissue Location

Highly expressed in placenta, spleen and peripheral blood leukocytes (PubMed:9435236, PubMed:9237759). Detected in monocytes, macrophages, dendritic cells and several types of T-cells (PubMed:9237759, PubMed:27022195). Expressed in pancreatic cancer cells but not in normal pancreatic cells (at protein level) (PubMed:36232715).

### Anti-TLR4 Antibody (aa420-435) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-TLR4 Antibody (aa420-435) - Images**