

Anti-SYNCRIP / HnRNP Q Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17959**Specification**

Anti-SYNCRIP / HnRNP Q Antibody - Product Information

Application	WB, IHC-P, E
Primary Accession	O60506
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	69603

Anti-SYNCRIP / HnRNP Q Antibody - Additional Information**Gene ID** 10492**Alias Symbol** **SYNCRIP****Other Names**

SYNCRIP, DJ3J17.2, GRYRBP, HnRNP-Q, HNRPO, HNRPO1, HnRNP Q, NS1-associated protein 1, NSAP1, GRY-RBP, PP68

Target/Specificity

Human SYNCRIP / HnRNP Q

Reconstitution & Storage

Caprylic acid and ammonium sulfate precipitation

Precautions

Anti-SYNCRIP / HnRNP Q Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-SYNCRIP / HnRNP Q Antibody - Protein Information**Name** SYNCRIP**Synonyms** HNRPO, NSAP1**Function**

Heterogenous nuclear ribonucleoprotein (hnRNP) implicated in mRNA processing mechanisms. Component of the CRD-mediated complex that promotes MYC mRNA stability. Isoform 1, isoform 2 and isoform 3 are associated in vitro with pre-mRNA, splicing intermediates and mature mRNA protein complexes. Isoform 1 binds to apoB mRNA AU-rich sequences. Isoform 1 is part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay

interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. Isoform 3 may be involved in cytoplasmic vesicle-based mRNA transport through interaction with synaptotagmins. Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma activation assembles into the GAIT complex which binds to stem loop- containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation; seems not to be essential for GAIT complex function.

Cellular Location

Cytoplasm. Microsome {ECO:0000250|UniProtKB:Q7TMK9} Endoplasmic reticulum. Nucleus {ECO:0000250|UniProtKB:Q7TMK9}. Note=The tyrosine phosphorylated form bound to RNA is found in microsomes (By similarity). Localized in cytoplasmic mRNP granules containing untranslated mRNAs (By similarity). {ECO:0000250|UniProtKB:O43390, ECO:0000250|UniProtKB:Q7TMK9} [Isoform 2]: Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q7TMK9}. Note=Expressed predominantly in the nucleoplasm. {ECO:0000250|UniProtKB:Q7TMK9}

Tissue Location

Ubiquitously expressed. Detected in heart, brain, pancreas, placenta, spleen, lung, liver, skeletal muscle, kidney, thymus, prostate, uterus, small intestine, colon, peripheral blood and testis.

Anti-SYNCRIP / HnRNP Q Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SYNCRIP / HnRNP Q Antibody - Images