

Anti-YBX1 / YB1 Antibody (clone 4F12)
Mouse Anti Human Monoclonal Antibody
Catalog # ALS17987

Specification

Anti-YBX1 / YB1 Antibody (clone 4F12) - Product Information

Application	WB, IHC-P, IF, E
Primary Accession	P67809
Predicted	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a,k
Calculated MW	35924

Anti-YBX1 / YB1 Antibody (clone 4F12) - Additional Information

Gene ID 4904

Alias Symbol YBX1

Other Names

YBX1, CBF-A, DBPB, CSDB, DNA-binding protein B, MDR-NF1, NSEP-1, NSEP1, Y box binding protein 1, Y-box transcription factor, Y-box-binding protein 1, YB-1, BP-8, CSDA2, EFI-A, Enhancer factor I subunit A, YB1

Target/Specificity

Human YBX1 / YB1

Reconstitution & Storage

Protein A purified

Precautions

Anti-YBX1 / YB1 Antibody (clone 4F12) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-YBX1 / YB1 Antibody (clone 4F12) - Protein Information

Name YBX1 ([HGNC:8014](#))

Function

DNA- and RNA-binding protein involved in various processes, such as translational repression, RNA stabilization, mRNA splicing, DNA repair and transcription regulation (PubMed: 8188694, PubMed: 10817758, PubMed: 11698476, PubMed: 14718551, PubMed: 18809583, PubMed: 31358969).

Predominantly acts as a RNA-binding protein: binds preferentially to the 5'-[CU]CUGCG-3' RNA

motif and specifically recognizes mRNA transcripts modified by C5-methylcytosine (m5C) (PubMed:19561594, PubMed:31358969). Promotes mRNA stabilization: acts by binding to m5C- containing mRNAs and recruiting the mRNA stability maintainer ELAVL1, thereby preventing mRNA decay (PubMed:10817758, PubMed:11698476, PubMed:31358969). Component of the CRD-mediated complex that promotes MYC mRNA stability (PubMed:19029303). Contributes to the regulation of translation by modulating the interaction between the mRNA and eukaryotic initiation factors (By similarity). Plays a key role in RNA composition of extracellular exosomes by defining the sorting of small non-coding RNAs, such as tRNAs, Y RNAs, Vault RNAs and miRNAs (PubMed:27559612, PubMed:29073095). Probably sorts RNAs in exosomes by recognizing and binding C5-methylcytosine (m5C)-containing RNAs (PubMed:28341602, PubMed:29073095). Acts as a key effector of epidermal progenitors by preventing epidermal progenitor senescence: acts by regulating the translation of a senescence-associated subset of cytokine mRNAs, possibly by binding to m5C-containing mRNAs (PubMed:29712925). Also involved in pre-mRNA alternative splicing regulation: binds to splice sites in pre-mRNA and regulates splice site selection (PubMed:12604611). Binds to TSC22D1 transcripts, thereby inhibiting their translation and negatively regulating TGF-beta- mediated transcription of COL1A2 (By similarity). Also able to bind DNA: regulates transcription of the multidrug resistance gene MDR1 is enhanced in presence of the APEX1 acetylated form at 'Lys-6' and 'Lys- 7' (PubMed:18809583). Binds to promoters that contain a Y-box (5'- CTGATTGCCAA-3'), such as MDR1 and HLA class II genes (PubMed:8188694, PubMed:18809583). Promotes separation of DNA strands that contain mismatches or are modified by cisplatin (PubMed:14718551). Has endonucleolytic activity and can introduce nicks or breaks into double- stranded DNA, suggesting a role in DNA repair (PubMed:14718551). The secreted form acts as an extracellular mitogen and stimulates cell migration and proliferation (PubMed:19483673).

Cellular Location

Cytoplasm. Nucleus. Cytoplasmic granule. Secreted. Secreted, extracellular exosome. Cytoplasm, P-body {ECO:0000250|UniProtKB:P62960}. Note=Predominantly cytoplasmic in proliferating cells (PubMed:12604611). Cytotoxic stress and DNA damage enhance translocation to the nucleus (PubMed:14718551) Localized in cytoplasmic mRNP granules containing untranslated mRNAs (PubMed:25229427). Shuttles between nucleus and cytoplasm (PubMed:25229427). Localized with DDX1, MBNL1 and TIAL1 in stress granules upon stress (PubMed:18335541). Secreted by mesangial and monocytic cells after inflammatory challenges (PubMed:19483673)

Anti-YBX1 / YB1 Antibody (clone 4F12) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-YBX1 / YB1 Antibody (clone 4F12) - Images