

**Anti-KRT19 / CK19 / Cytokeratin 19 Antibody (aa300-349)**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS18093****Specification**

---

**Anti-KRT19 / CK19 / Cytokeratin 19 Antibody (aa300-349) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P, E           |
| Primary Accession | <a href="#">P08727</a> |
| Predicted         | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | IgG                    |
| Calculated MW     | 44106                  |

**Anti-KRT19 / CK19 / Cytokeratin 19 Antibody (aa300-349) - Additional Information****Gene ID** 3880**Alias Symbol** **KRT19****Other Names**

KRT19, CK19, Cytokeratin 19, Cytokeratin-19, Keratin 19, Keratin-19, CK-19, K19, K1CS, Keratin, type I, 40-kd

**Target/Specificity**

Cytokeratin19 antibody detects endogenous levels of Cytokeratin19.

**Reconstitution & Storage**

Immunoaffinity purified

**Precautions**

Anti-KRT19 / CK19 / Cytokeratin 19 Antibody (aa300-349) is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-KRT19 / CK19 / Cytokeratin 19 Antibody (aa300-349) - Protein Information****Name** KRT19**Function**

Involved in the organization of myofibers. Together with KRT8, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle.

**Tissue Location**

Expressed in a defined zone of basal keratinocytes in the deep outer root sheath of hair follicles. Also observed in sweat gland and mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ectocervical epithelium (at protein level). Expressed in epidermal basal cells, in nipple epidermis and a defined region of the hair follicle. Also seen in a subset of vascular wall cells in both the veins and artery of human umbilical cord, and in umbilical cord vascular smooth muscle. Observed in muscle fibers

accumulating in the costameres of myoplasm at the sarcolemma in structures that contain dystrophin and spectrin.

### **Anti-KRT19 / CK19 / Cytokeratin 19 Antibody (aa300-349) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-KRT19 / CK19 / Cytokeratin 19 Antibody (aa300-349) - Images**