

Anti-SETD2 Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18117**Specification**

Anti-SETD2 Antibody - Product Information

Application	WB, IHC-P, IF, IP, CHIP
Primary Accession	Q9BYW2
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	287597

Anti-SETD2 Antibody - Additional Information**Gene ID** 29072

Alias Symbol	SETD2
--------------	--------------

Other Names

SETD2, HBP231, HIF-1, HIP-1, HSPC069, KMT3A, Lysine N-methyltransferase 3A, HIF1, HSET2, HYPB, p231HBP, SET2, Huntingtin yeast partner B, KIAA1732, SET domain containing 2

Target/Specificity

Human SETD2

Reconstitution & Storage

Immunoaffinity purified

Precautions

Anti-SETD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-SETD2 Antibody - Protein Information**Name** SETD2**Function**

Histone methyltransferase that specifically trimethylates 'Lys-36' of histone H3 (H3K36me3) using dimethylated 'Lys-36' (H3K36me2) as substrate (PubMed:[16118227](http://www.uniprot.org/citations/16118227), PubMed:[19141475](http://www.uniprot.org/citations/19141475), PubMed:[21526191](http://www.uniprot.org/citations/21526191), PubMed:[21792193](http://www.uniprot.org/citations/21792193), PubMed:[23043551](http://www.uniprot.org/citations/23043551), PubMed:[27474439](http://www.uniprot.org/citations/27474439)). It is capable of trimethylating unmethylated H3K36 (H3K36me0) in vitro (PubMed:[19332550](http://www.uniprot.org/citations/19332550)). Represents the main enzyme generating H3K36me3, a specific tag for epigenetic transcriptional activation (By

similarity). Plays a role in chromatin structure modulation during elongation by coordinating recruitment of the FACT complex and by interacting with hyperphosphorylated POLR2A (PubMed:23325844). Acts as a key regulator of DNA mismatch repair in G1 and early S phase by generating H3K36me3, a mark required to recruit MSH6 subunit of the MutS alpha complex: early recruitment of the MutS alpha complex to chromatin to be replicated allows a quick identification of mismatch DNA to initiate the mismatch repair reaction (PubMed:23622243). Required for DNA double-strand break repair in response to DNA damage: acts by mediating formation of H3K36me3, promoting recruitment of RAD51 and DNA repair via homologous recombination (HR) (PubMed:24843002). Acts as a tumor suppressor (PubMed:24509477). H3K36me3 also plays an essential role in the maintenance of a heterochromatic state, by recruiting DNA methyltransferase DNMT3A (PubMed:27317772). H3K36me3 is also enhanced in intron-containing genes, suggesting that SETD2 recruitment is enhanced by splicing and that splicing is coupled to recruitment of elongating RNA polymerase (PubMed:21792193). Required during angiogenesis (By similarity). Required for endoderm development by promoting embryonic stem cell differentiation toward endoderm: acts by mediating formation of H3K36me3 in distal promoter regions of FGFR3, leading to regulate transcription initiation of FGFR3 (By similarity). In addition to histones, also mediates methylation of other proteins, such as tubulins and STAT1 (PubMed:27518565, PubMed:28753426). Trimethylates 'Lys-40' of alpha-tubulins such as TUBA1B (alpha-TubK40me3); alpha-TubK40me3 is required for normal mitosis and cytokinesis and may be a specific tag in cytoskeletal remodeling (PubMed:27518565). Involved in interferon-alpha-induced antiviral defense by mediating both monomethylation of STAT1 at 'Lys-525' and catalyzing H3K36me3 on promoters of some interferon-stimulated genes (ISGs) to activate gene transcription (PubMed:28753426).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:E9Q5F9}. Chromosome {ECO:0000250|UniProtKB:E9Q5F9}

Tissue Location

Ubiquitously expressed.

Anti-SETD2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-SETD2 Antibody - Images