

**Anti-FGR Antibody (clone 3G10)**  
**Mouse Anti Human Monoclonal Antibody**  
**Catalog # ALS18150****Specification**

---

**Anti-FGR Antibody (clone 3G10) - Product Information**

Application	WB, IHC-P, E
Primary Accession	<a href="#">P09769</a>
Predicted	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a,k
Calculated MW	59479

**Anti-FGR Antibody (clone 3G10) - Additional Information****Gene ID** 2268**Alias Symbol** **FGR****Other Names**

FGR, C-fgr protooncogene, C-src2, C-fgr, Fgr kinase, Fgr oncogene, p55-Fgr, p58-Fgr, p58c-fgr, Proto-oncogene c-Fgr, Tyrosine-protein kinase Fgr, p55c-fgr, SRC2, C-src-2 proto-oncogene, Fgr proto oncogene, p55-c-fgr protein

**Target/Specificity**

Human FGR

**Reconstitution & Storage**

Protein A purified

**Precautions**

Anti-FGR Antibody (clone 3G10) is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-FGR Antibody (clone 3G10) - Protein Information****Name** FGR**Synonyms** SRC2**Function**

Non-receptor tyrosine-protein kinase that transmits signals from cell surface receptors devoid of kinase activity and contributes to the regulation of immune responses, including neutrophil, monocyte, macrophage and mast cell functions, cytoskeleton remodeling in response to extracellular stimuli, phagocytosis, cell adhesion and migration. Promotes mast cell degranulation, release of inflammatory cytokines and IgE-mediated anaphylaxis. Acts downstream of receptors that bind the Fc region of immunoglobulins, such as MS4A2/FCER1B, FCGR2A and/or FCGR2B. Acts downstream of ITGB1 and ITGB2, and regulates actin cytoskeleton reorganization, cell spreading

and adhesion. Depending on the context, activates or inhibits cellular responses. Functions as a negative regulator of ITGB2 signaling, phagocytosis and SYK activity in monocytes. Required for normal ITGB1 and ITGB2 signaling, normal cell spreading and adhesion in neutrophils and macrophages. Functions as a positive regulator of cell migration and regulates cytoskeleton reorganization via RAC1 activation. Phosphorylates SYK (in vitro) and promotes SYK-dependent activation of AKT1 and MAP kinase signaling. Phosphorylates PLD2 in antigen-stimulated mast cells, leading to PLD2 activation and the production of the signaling molecules lysophosphatidic acid and diacylglycerol. Promotes activation of PIK3R1. Phosphorylates FASLG, and thereby regulates its ubiquitination and subsequent internalization. Phosphorylates ABL1. Promotes phosphorylation of CBL, CTTN, PIK3R1, PTK2/FAK1, PTK2B/PYK2 and VAV2. Phosphorylates HCLS1 that has already been phosphorylated by SYK, but not unphosphorylated HCLS1. Together with CLNK, it acts as a negative regulator of natural killer cell-activating receptors and inhibits interferon-gamma production (By similarity).

#### **Cellular Location**

Cell membrane; Lipid-anchor; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle membrane. Cytoplasm, cytosol. Cytoplasm, cytoskeleton. Mitochondrion inner membrane. Mitochondrion intermembrane space. Note=Detected in mitochondrial intermembrane space and at inner membranes (By similarity). Colocalizes with actin fibers at membrane ruffles. Detected at plasma membrane lipid rafts

#### **Tissue Location**

Detected in neutrophils, monocytes and natural killer cells (at protein level). Detected in monocytes and large lymphocytes.

### **Anti-FGR Antibody (clone 3G10) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-FGR Antibody (clone 3G10) - Images**