

**Anti-CDC42 Antibody**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS18161****Specification**

---

**Anti-CDC42 Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">P60953</a>
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	21259

**Anti-CDC42 Antibody - Additional Information****Gene ID 998**Alias Symbol **CDC42****Other Names**

CDC42, G25K, G25K GTP-binding protein, Growth-regulating protein, CDC42Hs, GTP-binding protein, 25kD

**Target/Specificity**

Human CDC42

**Reconstitution & Storage**

Affinity purified

**Precautions**

Anti-CDC42 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-CDC42 Antibody - Protein Information****Name** CDC42 ([HGNC:1736](#))**Function**

Plasma membrane-associated small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses. Involved in epithelial cell polarization processes. Regulates the bipolar attachment of spindle microtubules to kinetochores before chromosome congression in metaphase (PubMed:<a href="http://www.uniprot.org/citations/15642749" target="\_blank">15642749</a>). Regulates cell migration (PubMed:<a href="http://www.uniprot.org/citations/17038317" target="\_blank">17038317</a>). In neurons, plays a role in the extension and maintenance of the formation of filopodia, thin and actin-rich surface projections (PubMed:<a href="http://www.uniprot.org/citations/14978216" target="\_blank">14978216</a>). Required for DOCK10-mediated spine formation in Purkinje cells and hippocampal neurons. In podocytes,

facilitates filopodia and podosomes formation upon DOCK11- activation (PubMed:<a href="http://www.uniprot.org/citations/33523862" target="\_blank">33523862</a>). Upon activation by CaMKII, modulates dendritic spine structural plasticity by relaying CaMKII transient activation to synapse-specific, long-term signaling (By similarity). Also plays a role in phagocytosis through organization of the F-actin cytoskeleton associated with forming phagocytic cups (PubMed:<a href="http://www.uniprot.org/citations/26465210" target="\_blank">26465210</a>).

#### **Cellular Location**

Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Midbody Cell projection, dendrite {ECO:0000250|UniProtKB:P60766} Note=Localizes to spindle during prometaphase cells. Moves to the central spindle as cells progressed through anaphase to telophase (PubMed:15642749). Localizes at the end of cytokinesis in the intercellular bridge formed between two daughter cells (PubMed:15642749). Its localization is regulated by the activities of guanine nucleotide exchange factor ECT2 and GTPase activating protein RACGAP1 (PubMed:15642749). Colocalizes with NEK6 in the centrosome (PubMed:20873783). In its active GTP-bound form localizes to the leading edge membrane of migrating dendritic cells (By similarity) {ECO:0000250|UniProtKB:P60766, ECO:0000269|PubMed:15642749, ECO:0000269|PubMed:20873783}

#### **Anti-CDC42 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-CDC42 Antibody - Images**