

#### **Anti-HNRNPA2B1 Antibody**

Rabbit Anti Human Polyclonal Antibody Catalog # ALS18174

## **Specification**

# **Anti-HNRNPA2B1 Antibody - Product Information**

Application WB, IHC-P, IF

Primary Accession <u>P22626</u>

Predicted Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 37430

# **Anti-HNRNPA2B1 Antibody - Additional Information**

**Gene ID 3181** 

Alias Symbol HNRNPA2B1

**Other Names** 

HNRNPA2B1, HNRNPB1, HNRPA2, HNRPA2B1, HNRPB1, HnRNP A2 / hnRNP B1, SNRPB1, HnRNP A2/B1, HNRNPA2, RNPA2

Target/Specificity Human HNRNPA2B1

Reconstitution & Storage Immunoaffinity purified

## **Precautions**

Anti-HNRNPA2B1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### **Anti-HNRNPA2B1 Antibody - Protein Information**

Name HNRNPA2B1

Synonyms HNRPA2B1

#### **Function**

Heterogeneous nuclear ribonucleoprotein (hnRNP) that associates with nascent pre-mRNAs, packaging them into hnRNP particles. The hnRNP particle arrangement on nascent hnRNA is non-random and sequence-dependent and serves to condense and stabilize the transcripts and minimize tangling and knotting. Packaging plays a role in various processes such as transcription, pre-mRNA processing, RNA nuclear export, subcellular location, mRNA translation and stability of mature mRNAs (PubMed:<a href="http://www.uniprot.org/citations/19099192" target="\_blank">19099192</a>). Forms hnRNP particles with at least 20 other different hnRNP and heterogeneous nuclear RNA in the nucleus. Involved in transport of specific mRNAs to the cytoplasm in oligodendrocytes and neurons: acts by specifically recognizing and binding the A2RE





(21 nucleotide hnRNP A2 response element) or the A2RE11 (derivative 11 nucleotide oligonucleotide) sequence motifs present on some mRNAs, and promotes their transport to the cytoplasm (PubMed:<a href="http://www.uniprot.org/citations/10567417"

target=" blank">10567417</a>). Specifically binds single-stranded telomeric DNA sequences, protecting telomeric DNA repeat against endonuclease digestion (By similarity). Also binds other RNA molecules, such as primary miRNA (pri-miRNAs): acts as a nuclear 'reader' of the N6-methyladenosine (m6A) mark by specifically recognizing and binding a subset of nuclear m6A-containing pri-miRNAs. Binding to m6A-containing pri-miRNAs promotes pri-miRNA processing by enhancing binding of DGCR8 to pri-miRNA transcripts (PubMed:<a href="http://www.uniprot.org/citations/26321680" target="\_blank">26321680</a>). Involved in miRNA sorting into exosomes following sumoylation, possibly by binding (m6A)-containing pre-miRNAs (PubMed: <a href="http://www.uniprot.org/citations/24356509" target=" blank">24356509</a>). Acts as a regulator of efficiency of mRNA splicing, possibly by

binding to m6A-containing pre-mRNAs (PubMed:<a

href="http://www.uniprot.org/citations/26321680" target=" blank">26321680</a>). Plays a role in the splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:<a href="http://www.uniprot.org/citations/20010808" target=" blank">20010808</a>). Also plays a role in the activation of the innate immune response (PubMed: <a href="http://www.uniprot.org/citations/31320558" target=" blank">31320558</a>). Mechanistically, senses the presence of viral DNA in the nucleus, homodimerizes and is demethylated by IMID6 (PubMed:<a href="http://www.uniprot.org/citations/31320558" target=" blank">31320558</a>). In turn, translocates to the cytoplasm where it activates the TBK1-IRF3 pathway, leading to interferon alpha/beta production (PubMed: <a href="http://www.uniprot.org/citations/31320558" target=" blank">31320558</a>).

#### **Cellular Location**

Nucleus. Nucleus, nucleoplasm. Cytoplasm. Cytoplasmic granule. Secreted, extracellular exosome. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs (PubMed:17289661). Component of ribonucleosomes (PubMed:17289661). Not found in the nucleolus (PubMed:17289661). Found in exosomes following sumoylation (PubMed:24356509).

# **Anti-HNRNPA2B1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cvtometv
- Cell Culture

Anti-HNRNPA2B1 Antibody - Images