

**Anti-AQP4 / Aquaporin 4 Antibody**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS18335****Specification**

---

**Anti-AQP4 / Aquaporin 4 Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IHC-P              |
| Primary Accession | <a href="#">P55087</a> |
| Predicted         | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | IgG                    |
| Calculated MW     | 34830                  |

**Anti-AQP4 / Aquaporin 4 Antibody - Additional Information****Gene ID 361**

|                                                                            |      |
|----------------------------------------------------------------------------|------|
| Alias Symbol                                                               | AQP4 |
| <b>Other Names</b>                                                         |      |
| AQP4, Aquaporin type4, Aquaporin 4, Aquaporin-4, HMIWC2, MIWC, WCH4, AQP-4 |      |

**Target/Specificity**

Human AQP4 / Aquaporin 4

**Reconstitution & Storage**

Affinity purified

**Precautions**

Anti-AQP4 / Aquaporin 4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-AQP4 / Aquaporin 4 Antibody - Protein Information****Name AQP4****Function**

Forms a water-specific channel (PubMed:<<http://www.uniprot.org/citations/7559426>>7559426</a>, PubMed:<<http://www.uniprot.org/citations/8601457>>8601457</a>, PubMed:<<http://www.uniprot.org/citations/19383790>>19383790</a>). Plays an important role in brain water homeostasis (PubMed:<<http://www.uniprot.org/citations/37143309>>37143309</a>). It is involved in glymphatic solute transport and is required for a normal rate of water exchange across the blood brain interface. Required for normal levels of cerebrospinal fluid influx into the brain cortex and parenchyma along paravascular spaces that surround penetrating arteries, and for normal drainage of interstitial fluid along paravenous drainage pathways. Thereby, it is required for normal clearance of solutes from the brain interstitial fluid, including soluble beta-amyloid peptides derived from APP. Plays a redundant role in urinary water homeostasis and urinary

concentrating ability (By similarity).

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Basolateral cell membrane {ECO:0000250|UniProtKB:P55088}; Multi-pass membrane protein. Endosome membrane {ECO:0000250|UniProtKB:P47863}. Cell membrane, sarcolemma; Multi-pass membrane protein. Cell projection {ECO:0000250|UniProtKB:P47863}. Note=Activation of the vasopressin receptor AVPR1A triggers AQP4 phosphorylation at Ser-180 and promotes its internalization from the cell membrane. Detected on brain astrocyte processes and astrocyte endfeet close to capillaries {ECO:0000250|UniProtKB:P47863}

#### **Tissue Location**

Detected in skeletal muscle (PubMed:29055082). Detected in stomach, along the glandular base region of the fundic gland (at protein level) (PubMed:8601457). Detected in brain, lung and skeletal muscle, and at much lower levels in heart and ovary (PubMed:7559426, PubMed:8601457).

### **Anti-AQP4 / Aquaporin 4 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-AQP4 / Aquaporin 4 Antibody - Images**