

Anti-Apelin Antibody (C-Terminus)

Rabbit Anti Human Polyclonal Antibody Catalog # ALS18454

Specification

Anti-Apelin Antibody (C-Terminus) - Product Information

Application WB, IHC-P Primary Accession O9ULZ1

Predicted Human, Mouse, Rat, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 8569

Anti-Apelin Antibody (C-Terminus) - Additional Information

Gene ID 8862

Alias Symbol APLN

Other Names

APLN, AGTRL1 ligand, Apelin, APEL, Preproapelin, APJ endogenous ligand, XNPEP2

Target/Specificity

Recognizes endogenous levels of Apelin protein.

Reconstitution & Storage

Immunoaffinity purified

Precautions

Anti-Apelin Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Apelin Antibody (C-Terminus) - Protein Information

Name APLN

Synonyms APEL

Function

Endogenous ligand for the apelin receptor (APLNR) (PubMed: 10525157). Drives internalization of the apelin receptor (By similarity). Apelin-36 dissociates more hardly than (pyroglu)apelin-13 from APLNR (By similarity). Hormone involved in the regulation of cardiac precursor cell movements during gastrulation and heart morphogenesis (By similarity). Has an inhibitory effect on cytokine production in response to T-cell receptor/CD3 cross-linking; the oral intake of apelin in the colostrum and the milk might therefore modulate immune responses in neonates (By similarity). Plays a role in early coronary blood vessels formation (By similarity). Mediates myocardial contractility in an ERK1/2-dependent manner (By similarity). May also have a role in the central control of body fluid homeostasis by influencing vasopressin release and



drinking behavior (By similarity).

Cellular Location

Secreted {ECO:0000250|UniProtKB:Q9TUI9}. Secreted, extracellular space {ECO:0000250|UniProtKB:Q9TUI9} Note=Abundantly secreted in the colostrum. Lower level in milk Decreases rapidly within several days after parturition in milk, but is still detectable even in commercial milk {ECO:0000250|UniProtKB:Q9TUI9}

Tissue Location

Expressed in the brain with highest levels in the frontal cortex, thalamus, hypothalamus and midbrain (PubMed:10617103) Secreted by the mammary gland into the colostrum and the milk

Anti-Apelin Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Apelin Antibody (C-Terminus) - Images