

**Anti-CCL24 / Eotaxin 2 Antibody (aa27-119)**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS18549****Specification**

---

**Anti-CCL24 / Eotaxin 2 Antibody (aa27-119) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC, IHC-P, E          |
| Primary Accession | <a href="#">O00175</a> |
| Predicted         | Human                  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Isotype           | IgG                    |
| Calculated MW     | 13134                  |

**Anti-CCL24 / Eotaxin 2 Antibody (aa27-119) - Additional Information****Gene ID** 6369**Alias Symbol** CCL24**Other Names**

CCL24, C-C motif chemokine 24, Ckb-6, CK-beta-6, Eotaxin2, MPIF-2, MPIF2, SCYA24, Small-inducible cytokine A24, Eotaxin-2

**Target/Specificity**

Human CCL24 / Eotax 2

**Reconstitution & Storage**

Caprylic acid and ammonium sulfate precipitation

**Precautions**

Anti-CCL24 / Eotaxin 2 Antibody (aa27-119) is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-CCL24 / Eotaxin 2 Antibody (aa27-119) - Protein Information****Name** CCL24 ([HGNC:10623](#))**Function**

Chemotactic for resting T-lymphocytes, and eosinophils (PubMed:<a href="http://www.uniprot.org/citations/9104803" target="\_blank">9104803</a>, PubMed:<a href="http://www.uniprot.org/citations/9365122" target="\_blank">9365122</a>). Has lower chemotactic activity for neutrophils but none for monocytes and activated lymphocytes (PubMed:<a href="http://www.uniprot.org/citations/9104803" target="\_blank">9104803</a>, PubMed:<a href="http://www.uniprot.org/citations/9365122" target="\_blank">9365122</a>). Is a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line (PubMed:<a href="http://www.uniprot.org/citations/9104803" target="\_blank">9104803</a>, PubMed:<a href="http://www.uniprot.org/citations/9365122" target="\_blank">9365122</a>). Binds to CCR3 (PubMed:<a href="http://www.uniprot.org/citations/9104803" target="\_blank">9104803</a>)

target="\_blank">9104803</a>, PubMed:<a href="http://www.uniprot.org/citations/9365122" target="\_blank">9365122</a>).

**Cellular Location**

Secreted

**Tissue Location**

Activated monocytes and activated T lymphocytes.

**Anti-CCL24 / Eotaxin 2 Antibody (aa27-119) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-CCL24 / Eotaxin 2 Antibody (aa27-119) - Images**