

**GSTA1 Antibody**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM1932b****Specification**

---

**GSTA1 Antibody - Product Information**

Application	WB,E
Primary Accession	<a href="#">P08263</a>
Other Accession	<a href="#">NP_665683.1</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgM,k

**GSTA1 Antibody - Additional Information****Gene ID** 2938**Other Names**

Glutathione S-transferase A1, GST HA subunit 1, GST class-alpha member 1, GST-epsilon, GSTA1-1, GTH1, Glutathione S-transferase A1, N-terminally processed, GSTA1

**Target/Specificity**

This GSTA1 monoclonal antibody is generated from mouse immunized with GSTA1 recombinant protein.

**Dilution**

WB~~1:100

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GSTA1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**GSTA1 Antibody - Protein Information****Name** GSTA1

**Function** Glutathione S-transferase that catalyzes the nucleophilic attack of the sulfur atom of glutathione on the electrophilic groups of a wide range of exogenous and endogenous compounds (Probable). Involved in the formation of glutathione conjugates of both prostaglandin A2 (PGA2) and prostaglandin J2 (PGJ2) (PubMed:[9084911](#)). It also catalyzes the isomerization of

D5-androstene-3,17-dione (AD) into D4-androstene- 3,17-dione and may therefore play an important role in hormone biosynthesis (PubMed:[11152686](#)). Through its glutathione-dependent peroxidase activity toward the fatty acid hydroperoxide (13S)-hydroperoxy-(9Z,11E)-octadecadienoate/13-HPODE it is also involved in the metabolism of oxidized linoleic acid (PubMed:[16624487](#)).

**Cellular Location**

Cytoplasm.


**Tissue Location**

Liver.

**GSTA1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**GSTA1 Antibody - Images**

GSTA1 Antibody (Cat. #AM1932b) western blot analysis in HepG2 cell line lysates (35µg/lane). This demonstrates the GSTA1 antibody detected the GSTA1 protein (arrow).

**GSTA1 Antibody - Background**

Cytosolic and membrane-bound forms of glutathione S-transferase are encoded by two distinct supergene families. These enzymes function in the detoxification of electrophilic compounds, including carcinogens, therapeutic drugs, environmental toxins and products of oxidative stress, by conjugation with glutathione. The genes encoding these enzymes are known to be highly polymorphic. These genetic variations can change an individual's susceptibility to carcinogens and toxins as well as affect the toxicity and

efficacy of some drugs. At present, eight distinct classes of the soluble cytoplasmic mammalian glutathione S-transferases have been identified: alpha, kappa, mu, omega, pi, sigma, theta and zeta. This gene encodes a glutathione S-transferase belonging to the alpha class. The alpha class genes, located in a cluster mapped to chromosome 6, are the most abundantly expressed glutathione S-transferases in liver. In addition to metabolizing bilirubin and certain anti-cancer drugs in the liver, the alpha class of these enzymes exhibit glutathione peroxidase activity thereby protecting the cells from reactive oxygen species and the products of peroxidation.

#### **GSTA1 Antibody - References**

Elhasid, R., et al. *Pediatr Blood Cancer* 55(6):1172-1179(2010)  
Hawken, S.J., et al. *Hum. Genet.* 128(1):89-101(2010)  
Oguztuzun, S., et al. *Folia Histochem. Cytobiol.* 48(1):122-127(2010)  
Eriksen, K.T., et al. *J. Toxicol. Environ. Health Part A* 73(9):583-595(2010)  
Nguyen, T.V., et al. *Oncol. Res.* 18(7):349-355(2010)