

APCS Antibody (Ascites)

Mouse Monoclonal Antibody (Mab)
Catalog # AM1978a

Specification

APCS Antibody (Ascites) - Product Information

Application WB,E **Primary Accession** P02743 NP 001630.1 Other Accession Reactivity Human Host Mouse Clonality **Monoclonal** Isotype IgG2a Calculated MW 25387

APCS Antibody (Ascites) - Additional Information

Gene ID 325

Other Names

Serum amyloid P-component, SAP, 95S alpha-1-glycoprotein, Serum amyloid P-component(1-203), APCS, PTX2

Target/Specificity

Purified His-tagged APCS protein(Fragment) was used to produced this monoclonal antibody.

Dilution

WB~~1:500~4000

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

APCS Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

APCS Antibody (Ascites) - Protein Information

Name APCS

Synonyms PTX2

Function Can interact with DNA and histones and may scavenge nuclear material released from damaged circulating cells. May also function as a calcium-dependent lectin.



Cellular Location Secreted.


Tissue LocationFound in serum and urine.

APCS Antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

APCS Antibody (Ascites) - Images



APCS antibody (Cat. #AM1978a) western blot analysis in K562 cell line lysates ($35\mu g$ /lane). This demonstrates the APCS antibody detected the APCS protein (arrow).

APCS Antibody (Ascites) - Background

The protein encoded by this gene is a glycoprotein, belonging to the pentraxin family of proteins, which has a characteristic pentameric organization. These family members have considerable sequence homology which is thought to be the result of gene duplication. The binding of the encoded protein to proteins in the pathological amyloid cross-beta fold suggests its possible role as a chaperone. This protein is also thought to control the degradation of chromatin. It has been demonstrated that this protein binds to apoptotic cells at an early stage, which raises the possibility that it is involved in dealing with apoptotic cells in vivo.

APCS Antibody (Ascites) - References





Song, Z., et al. Atherosclerosis 211(1):90-95(2010) Davila, S., et al. Genes Immun. 11(3):232-238(2010)

McGeachie, M., et al. Circulation 120(24):2448-2454(2009)

Lu, J., et al. Nature 456(7224):989-992(2008)

Verwey, N.A., et al. Dement Geriatr Cogn Disord 26(6):522-527(2008)