

TNFSF4 Antibody (Ascites)
Mouse Monoclonal Antibody (Mab)
Catalog # AM1983a**Specification**

TNFSF4 Antibody (Ascites) - Product Information

Application	WB,E
Primary Accession	P23510
Other Accession	NP_003317.1
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Calculated MW	21050
Antigen Region	97-125

TNFSF4 Antibody (Ascites) - Additional Information**Gene ID** 7292**Other Names**

Tumor necrosis factor ligand superfamily member 4, Glycoprotein Gp34, OX40 ligand, OX40L, TAX transcriptionally-activated glycoprotein 1, CD252, TNFSF4, TXGP1

Target/Specificity

This TNFSF4 antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 97-125 amino acids from human TNFSF4.

Dilution

WB~~1:500~4000

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TNFSF4 Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

TNFSF4 Antibody (Ascites) - Protein Information**Name** TNFSF4**Synonyms** TXGP1

Function Cytokine that binds to TNFRSF4. Co-stimulates T-cell proliferation and cytokine production.

Cellular Location


Membrane; Single-pass type II membrane protein.

TNFSF4 Antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TNFSF4 Antibody (Ascites) - Images



TNFSF4 antibody (Cat. #AM1983a) western blot analysis in HL-60 cell line lysates (35µg/lane). This demonstrates the TNFSF4 antibody detected the TNFSF4 protein (arrow).

TNFSF4 Antibody (Ascites) - Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for receptor TNFRSF4/OX4. It is found to be involved in T cell antigen-presenting cell (APC) interactions. In surface Ig- and CD40-stimulated B cells, this cytokine along with CD70 has been shown to provide CD28-independent costimulatory signals to T cells. This protein and its receptor are reported to directly mediate adhesion of activated T cells to vascular endothelial cells.

TNFSF4 Antibody (Ascites) - References

Karulf, M., et al. J. Immunol. 185(8):4856-4862(2010)
Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Mosbruger, T.L., et al. J. Infect. Dis. 201(9):1371-1380(2010)
Davila, S., et al. Genes Immun. 11(3):232-238(2010)