

## **NUAK2 Antibody**

Mouse Monoclonal Antibody (Mab)
Catalog # AM1988b

## **Specification**

## **NUAK2 Antibody - Product Information**

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Calculated MW

WB,E Q9H093 NP\_112214.1 Human Mouse Monoclonal

IgM 69612

## **NUAK2 Antibody - Additional Information**

### **Gene ID 81788**

#### **Other Names**

NUAK family SNF1-like kinase 2, Omphalocele kinase 2, SNF1/AMP kinase-related kinase, SNARK, NUAK2 {ECO:0000312|EMBL:AAH173061}, OMPHK2, SNARK

### Target/Specificity

Purified His-tagged NUAK2 protein(Fragment) was used to produced this monoclonal antibody.

#### **Dilution**

WB~~1:500~1000

#### **Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.

# Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

NUAK2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **NUAK2 Antibody - Protein Information**

Name NUAK2 {ECO:0000312|EMBL:AAH17306.1}

Synonyms OMPHK2, SNARK

**Function** Stress-activated kinase involved in tolerance to glucose starvation. Induces cell-cell detachment by increasing F-actin conversion to G-actin. Expression is induced by CD95 or




TNF-alpha, via NF-kappa-B. Protects cells from CD95-mediated apoptosis and is required for the increased motility and invasiveness of CD95-activated tumor cells. Phosphorylates LATS1 and LATS2. Plays a key role in neural tube closure during embryonic development through LATS2 phosphorylation and regulation of the nuclear localization of YAP1 a critical downstream regulatory target in the Hippo signaling pathway (PubMed: 32845958).

## **NUAK2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **NUAK2 Antibody - Images**



NUAK2 Antibody (Cat. #AM1988b) western blot analysis in Hela cell line lysates (35μg/lane). This demonstrates the NUAK2 antibody detected the NUAK2 protein (arrow).

# **NUAK2 Antibody - Background**

Stress-activated kinase involved in tolerance to glucose starvation. Induces cell-cell detachment by increasing F-actin conversion to G-actin. Expression is induced by CD95 or TNF-alpha, via NF-kappa-B. Protects cells from CD95-mediated apoptosis and is required for the increased motility and invasiveness of CD95-activated tumor cells.

# **NUAK2 Antibody - References**

Koh, H.J., et al. Proc. Natl. Acad. Sci. U.S.A. 107(35):15541-15546(2010)

Rune, A., et al. Diabetologia 52(10):2182-2189(2009)

Kuga, W., et al. Biochem. Biophys. Res. Commun. 377(4):1062-1066(2008)

Kim, J.H., et al. Leuk. Lymphoma 49(7):1392-1398(2008)

Yamamoto, H., et al. Biochem. Biophys. Res. Commun. 365(3):541-547(2008)