

CD1A Antibody (Ascites)
Mouse Monoclonal Antibody (Mab)
Catalog # AM2043a**Specification**

CD1A Antibody (Ascites) - Product Information

Application	WB,E
Primary Accession	P06126
Other Accession	NP_001754.2
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	37077
Antigen Region	76-104

CD1A Antibody (Ascites) - Additional Information**Gene ID** 909**Other Names**

T-cell surface glycoprotein CD1a, T-cell surface antigen T6/Leu-6, hTa1 thymocyte antigen, CD1a, CD1A

Target/Specificity

This CD1A antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 76-104 amino acids from human CD1A.

Dilution

WB~~1:500~8000

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CD1A Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

CD1A Antibody (Ascites) - Protein Information**Name** CD1A

Function Antigen-presenting protein that binds self and non-self lipid and glycolipid antigens and presents them to T-cell receptors on natural killer T-cells.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Note=Subject to intracellular trafficking between the cell membrane and endosomes (PubMed:11231314). Localizes to cell surface lipid rafts (PubMed:18178838).


Tissue Location

Expressed on cortical thymocytes, epidermal Langerhans cells, dendritic cells, on certain T-cell leukemias, and in various other tissues.

CD1A Antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD1A Antibody (Ascites) - Images

CD1A Antibody (Cat. #AM2043a) western blot analysis in Jurkat cell line lysates (35µg/lane). This demonstrates the CD1A antibody detected the CD1A protein (arrow).

CD1A Antibody (Ascites) - Background

This gene encodes a member of the CD1 family of transmembrane glycoproteins, which are structurally related to the major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to the plasma membrane and to recycling

vesicles of the early endocytic system. Alternatively spliced transcript variants have been observed, but their biological validity has not been determined.

CD1A Antibody (Ascites) - References

Zeissig, S., et al. J. Clin. Invest. 120(8):2889-2899(2010)
Davila, S., et al. Genes Immun. 11(3):232-238(2010)
Valencia, J., et al. J. Leukoc. Biol. 87(3):405-414(2010)
Cernadas, M., et al. J. Immunol. 184(3):1235-1241(2010)
Young, D.C., et al. J. Biol. Chem. 284(37):25087-25096(2009)