

CYP3A4 Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM2065a**Specification**

CYP3A4 Antibody - Product Information

Application	WB,E
Primary Accession	P08684
Other Accession	NP_059488.2
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Antigen Region	228-255

CYP3A4 Antibody - Additional Information**Gene ID** 1576**Other Names**

Cytochrome P450 3A4, 11413-, 8-cineole 2-exo-monooxygenase, Albendazole monooxygenase, Albendazole sulfoxidase, CYP11A3, CYP11A4, Cytochrome P450 3A3, Cytochrome P450 HLp, Cytochrome P450 NF-25, Cytochrome P450-PCN1, Nifedipine oxidase, Quinine 3-monooxygenase, Taurochenodeoxycholate 6-alpha-hydroxylase, CYP3A4, CYP3A3

Target/Specificity

This CYP3A4 antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 228-255 amino acids from human CYP3A4.

Dilution

WB~~1:100~1000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CYP3A4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CYP3A4 Antibody - Protein Information**Name** CYP3A4 {ECO:0000303|PubMed:11470997, ECO:0000312|HGNC:HGNC:2637}**Function** A cytochrome P450 monooxygenase involved in the metabolism of sterols, steroid

hormones, retinoids and fatty acids (PubMed:[10681376](#), PubMed:[11093772](#), PubMed:[11555828](#), PubMed:[14559847](#), PubMed:[12865317](#), PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[20702771](#), PubMed:[19965576](#), PubMed:[21490593](#), PubMed:[21576599](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase). Catalyzes the hydroxylation of carbon-hydrogen bonds (PubMed:[2732228](#), PubMed:[14559847](#), PubMed:[12865317](#), PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[21576599](#), PubMed:[21490593](#)). Exhibits high catalytic activity for the formation of hydroxysteroids from estrone (E1) and 17beta- estradiol (E2), namely 2-hydroxy E1 and E2, as well as D-ring hydroxylated E1 and E2 at the C-16 position (PubMed:[11555828](#), PubMed:[14559847](#), PubMed:[12865317](#)). Plays a role in the metabolism of androgens, particularly in oxidative deactivation of testosterone (PubMed:[2732228](#), PubMed:[15373842](#), PubMed:[15764715](#), PubMed:[22773874](#)). Metabolizes testosterone to less biologically active 2beta- and 6beta- hydroxytestosterones (PubMed:[2732228](#), PubMed:[15373842](#), PubMed:[15764715](#)). Contributes to the formation of hydroxycholesterols (oxysterols), particularly A-ring hydroxylated cholesterol at the C- 4beta position, and side chain hydroxylated cholesterol at the C-25 position, likely contributing to cholesterol degradation and bile acid biosynthesis (PubMed:[21576599](#)). Catalyzes bisallylic hydroxylation of polyunsaturated fatty acids (PUFA) (PubMed:[9435160](#)). Catalyzes the epoxidation of double bonds of PUFA with a preference for the last double bond (PubMed:[19965576](#)). Metabolizes endocannabinoid arachidonoyl ethanolamide (anandamide) to 8,9-, 11,12-, and 14,15- epoxyeicosatrienoic acid ethanolamides (EpETRE-EAs), potentially modulating endocannabinoid system signaling (PubMed:[20702771](#)). Plays a role in the metabolism of retinoids. Displays high catalytic activity for oxidation of all-trans-retinol to all-trans-retinal, a rate- limiting step for the biosynthesis of all-trans-retinoic acid (atRA) (PubMed:[10681376](#)). Further metabolizes atRA toward 4-hydroxyretinoate and may play a role in hepatic atRA clearance (PubMed:[11093772](#)). Responsible for oxidative metabolism of xenobiotics. Acts as a 2-exo- monooxygenase for plant lipid 1,8-cineole (eucalyptol) (PubMed:[11159812](#)). Metabolizes the majority of the administered drugs. Catalyzes sulfoxidation of the anthelmintics albendazole and fenbendazole (PubMed:[10759686](#)). Hydroxylates antimalarial drug quinine (PubMed:[8968357](#)). Acts as a 1,4-cineole 2-exo-monoxygenase (PubMed:[11695850](#)). Also involved in vitamin D catabolism and calcium homeostasis. Catalyzes the inactivation of the active hormone calcitriol (1-alpha,25-dihydroxyvitamin D(3)) (PubMed:[29461981](#)).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein. Microsome membrane; Single-pass membrane protein

Tissue Location

Expressed in prostate and liver. According to some authors, it is not expressed in brain (PubMed:[19094056](#)). According to others, weak levels of expression are measured in some brain locations (PubMed:[19359404](#), PubMed:[18545703](#)). Also expressed in epithelium of the small intestine and large intestine, bile duct, nasal mucosa, kidney, adrenal cortex, epithelium of the gastric mucosa with intestinal metaplasia, gallbladder, intercalated ducts of the pancreas, chief cells of the parathyroid and the corpus luteum of the ovary (at protein level).


CYP3A4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

CYP3A4 Antibody - Images



CYP3A4 Antibody (Cat. #AM2065a) western blot analysis in T47D cell line lysates (35µg/lane). This demonstrates the CYP3A4 antibody detected the CYP3A4 protein (arrow).

CYP3A4 Antibody - Background

This gene, CYP3A4, encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by glucocorticoids and some pharmacological agents. This enzyme is involved in the metabolism of approximately half the drugs which are used today, including acetaminophen, codeine, cyclosporin A, diazepam and erythromycin. The enzyme also metabolizes some steroids and carcinogens. This gene is part of a cluster of cytochrome P450 genes on chromosome 7q21.1. Previously another CYP3A gene, CYP3A3, was thought to exist; however, it is now thought that this sequence represents a transcript variant of CYP3A4.

CYP3A4 Antibody - References

Justenhoven, C., et al. Cancer 116(23):5358-5364(2011)
Moore, C.D., et al. Biochemistry 49(41):9011-9019(2010)
Jablonski, K.A., et al. Diabetes 59(10):2672-2681(2010)
Hu, M., et al. Pharmacogenet. Genomics 20(10):634-637(2010)
Coto, E., et al. Biochem. Biophys. Res. Commun. (2010) In press :