

**CD33 Antibody**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM2149b****Specification**

---

**CD33 Antibody - Product Information**

Application	IF, WB,E
Primary Accession	<a href="#">P20138</a>
Other Accession	<a href="#">NP_001763.3</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Calculated MW	39825
Antigen Region	337-364

**CD33 Antibody - Additional Information****Gene ID** 945**Other Names**

Myeloid cell surface antigen CD33, Sialic acid-binding Ig-like lectin 3, Siglec-3, gp67, CD33, CD33, SIGLEC3

**Target/Specificity**

This CD33 antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 337-364 amino acids from human CD33.

**Dilution**

IF~~1:10~50  
WB~~1:500~1000

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CD33 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CD33 Antibody - Protein Information****Name** CD33**Synonyms** SIGLEC3

**Function** Sialic-acid-binding immunoglobulin-like lectin (Siglec) that plays a role in mediating cell-cell interactions and in maintaining immune cells in a resting state (PubMed:[10611343](#), PubMed:[15597323](#), PubMed:[11320212](#)). Preferentially recognizes and binds alpha-2,3- and more avidly alpha-2,6-linked sialic acid-bearing glycans (PubMed:[7718872](#)). Upon engagement of ligands such as C1q or sialylated glycoproteins, two immunoreceptor tyrosine-based inhibitory motifs (ITIMs) located in CD33 cytoplasmic tail are phosphorylated by Src-like kinases such as LCK (PubMed:[28325905](#), PubMed:[10887109](#)). These phosphorylations provide docking sites for the recruitment and activation of protein-tyrosine phosphatases PTPN6/SHP-1 and PTPN11/SHP-2 (PubMed:[10556798](#), PubMed:[10206955](#), PubMed:[10887109](#)). In turn, these phosphatases regulate downstream pathways through dephosphorylation of signaling molecules (PubMed:[10206955](#), PubMed:[10887109](#)). One of the repressive effects of CD33 on monocyte activation requires phosphoinositide 3-kinase/PI3K (PubMed:[15597323](#)).

#### Cellular Location

[Isoform CD33M]: Cell membrane; Single-pass type I membrane protein

#### Tissue Location


Monocytic/myeloid lineage cells. In the brain, CD33 is mainly expressed on microglial cells

### CD33 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD33 Antibody - Images



Fluorescent image of A549 cell stained with CD33 Antibody (Cat#AM2149b/SM120504AC). A549 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with CD33 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor®

488 conjugated donkey anti-mouse antibody (green) was used (1:400, 50 min at 37°C. Nuclei were counterstained with DAPI (blue) (10 µg/ml, 10 min).CD33 immunoreactivity is localized to Membrane significantly.



CD33 Antibody(Cat. #AM2149b) western blot analysis in HepG2 cell line lysates (35µg/lane).This demonstrates the CD33 antibody detected the CD33 protein (arrow).

### CD33 Antibody - Background

Putative adhesion molecule of myelomonocytic-derived cells that mediates sialic-acid dependent binding to cells. Preferentially binds to alpha-2,6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic acids on the same cell surface. In the immune response, may act as an inhibitory receptor upon ligand induced tyrosine phosphorylation by recruiting cytoplasmic phosphatase(s) via their SH2 domain(s) that block signal transduction through dephosphorylation of signaling molecules. Induces apoptosis in acute myeloid leukemia (in vitro).

### CD33 Antibody - References

- Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
- Davila, S., et al. Genes Immun. 11(3):232-238(2010)
- Akahane, K., et al. Leukemia 24(4):865-869(2010)
- Shamsasenjan, K., et al. Int. J. Hematol. 89(3):310-318(2009)
- Bertram, L., et al. Am. J. Hum. Genet. 83(5):623-632(2008)