

PAH Antibody (Center)
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8419b

Specification

PAH Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P00439
Other Accession	P04176 , P16331 , Q2KIH7
Reactivity	Human, Mouse, Rat
Predicted	Bovine
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, κ

PAH Antibody (Center) - Additional Information

Gene ID 5053

Other Names

Phenylalanine-4-hydroxylase, PAH, Phe-4-monooxygenase, PAH

Target/Specificity

This PAH antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 127-161 amino acids from the Central region of human PAH.

Dilution

WB~~1:1000

IHC-P~~1:25

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PAH Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PAH Antibody (Center) - Protein Information

Name PAH


Function Catalyzes the hydroxylation of L-phenylalanine to L-tyrosine.

PAH Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PAH Antibody (Center) - Images




Western blot analysis of lysates from HepG2 cell line and mouse liver rat liver tissue lysates (from left to right) using PAH Antibody (Center)(Cat. #AM8419b). AM8419b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.



Immunohistochemical analysis of paraffin-embedded H. liver section using PAH Antibody (Center)(Cat#AM8419b). AM8419b was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded M. kidney section using PAH Antibody (Center)(Cat#AM8419b). AM8419b was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



AM8419b staining PAH in human kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

PAH Antibody (Center) - References

Kwok S.C.M., et al. *Biochemistry* 24:556-561(1985).
Scriber C.R., et al. Submitted (SEP-1997) to the EMBL/GenBank/DDBJ databases.
Cotton R.G., et al. *Biochem. J.* 255:193-196(1988).
Miranda F.F., et al. *J. Biol. Chem.* 277:40937-40943(2002).
Siltberg-Liberles J., et al. *Gene* 427:86-92(2008).

PAH Antibody (Center) - Citations

- [Integrated Proteomics and Metabolomics Reveal the Mechanism of Nephrotoxicity Induced by Triptolide](#)