

Phospho-Thr292 MEK 1 Antibody
Affinity purified rabbit polyclonal antibody
Catalog # AN1016

Specification

Phospho-Thr292 MEK 1 Antibody - Product Information

Application	WB
Primary Accession	Q02750
Reactivity	Human
Predicted	Bovine, Human, Mouse, Monkey, Rat
Host	Rabbit
Clonality	polyclonal
Calculated MW	45 KDa

Phospho-Thr292 MEK 1 Antibody - Additional Information

Gene ID	5604
Gene Name	MAP2K1

Other Names

Dual specificity mitogen-activated protein kinase kinase 1, MAP kinase kinase 1, MAPKK 1, MKK1, ERK activator kinase 1, MAPK/ERK kinase 1, MEK 1, MAP2K1, MEK1, PRKMK1

Target/Specificity

Synthetic phospho-peptide corresponding to amino acid residues surrounding Thr292 conjugated to KLH.

Dilution

WB~~ 1:1000

Format

Prepared from rabbit serum by affinity purification via sequential chromatography on phospho- and dephosphopeptide affinity columns.

Antibody Specificity

Specific for the ~45k MEK 1 protein phosphorylated at Thr292 in Western blots.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-Thr292 MEK 1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Phospho-Thr292 MEK 1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-Thr292 MEK 1 Antibody - Images



Western blot of recombinant wild type and mutant MEK 1 showing immunolabeling of the ~45k MEK-1 protein phosphorylated at Thr292. Lanes 1 and 2 are WT MEK 1 and Lanes 3 and 4 are mutant MEK 1 (T292A). MAP kinase was coexpressed in the samples run in Lanes 2 and 4.

Phospho-Thr292 MEK 1 Antibody - Background

MEK 1 (MAP kinase kinase, also known as MKK) is an integral component of the MAP kinase cascade that regulates cell growth and differentiation (Ahn, 1993; Chong et al., 2003). This pathway also plays a key role in synaptic plasticity in the brain (Adams and Sweatt, 2002). Activated MEK 1 acts as a dual specificity kinase phosphorylating both a threonine and a tyrosine residue on MAP kinase (Kyriakis et al., 1991; Seger et al., 1991; Crews et al., 1992). Conversely, there also appears to be a feedback phosphorylation of MEK 1 by MAP kinase. The sites on MEK 1 that are phosphorylated by MAP kinase are Thr292 and Thr386 (Mansour et al., 1994).

Phospho-Thr292 MEK 1 Antibody - References

- Adams JP, Sweatt JD (2002) Molecular psychology: Roles for the ERK MAP kinase cascade in memory. *Annu Rev Pharmacol Toxicol* 42:135-163.
- Ahn NG (1993) The MAP kinase cascade. Discovery of a new signal transduction pathway. *Mol Cell Biochem* 127-128:201-209.
- Chong H, Vikis HG, Guan KL (2003) Mechanisms of regulating the Raf kinase family. *Cellular Signalling* 15:463-469.
- Crews CM, Alessandrini A, Erikson RL (1992) The primary structure of MEK, a protein kinase that phosphorylates the ERK gene product. *Science* 258:478-480.
- Kyriakis JM, Brautigan DL, Ingebritsen TS, Avruch J (1991) pp54 Microtubule-associated protein-2 kinase requires both tyrosine and serine/threonine phosphorylation for activity. *J Biol Chem* 266:10043-10046.
- Mansour SJ, Resing KA, Candi JM, Hermann AS, Gloor JW, Herskind KR, Wartmann M, Davis RJ, Ahn NG (1994) Mitogen-activated protein (MAP) kinase phosphorylation of MAP kinase kinase: Determination of phosphorylation sites by mass spectrometry and site-directed mutagenesis. *J Biochem (Tokyo)* 116:304-314.
- Seger R, Ahn NG, Boulton TG, Yancopoulos GD, Panayotatos N, Radziejewska E, Ericsson L, Bratlien

RL, Cobb MH, Krebs EG (1991) Microtubule-associated protein 2 kinases, ERK1 and ERK2, undergo autophosphorylation on both tyrosine and threonine residues: Implications for their mechanism of activation. Proc Natl Acad Sci USA 88:6142-6146.