10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CD3 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1024a

Specification

CD3 Antibody - Product Information

Application	**FC**
Primary Accession	P20963
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**IgG2a**

Description
CD3 is a component of T cell surface receptor (TCR), it cantains 5 invariable chains: gamma- , delta-, epsilon-, and zeta/eta□Clevers et al. (1988) showed the TCR/CD3 complex consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits□CD3 is responsible for the signal transduction of the TCR.

Immunogen
Purified recombinant fragment of human CD3 expressed in E. Coli.

Formulation
Ascitic fluid containing 0.03% sodium azide.

CD3 Antibody - Additional Information

Gene ID 919

Other Names
T-cell surface glycoprotein CD3 zeta chain, T-cell receptor T3 zeta chain, CD247, CD247, CD3Z, T3Z, TCRZ

Dilution
FC~~1/200 - 1/400

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CD3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CD3 Antibody - Protein Information

Name CD247

Synonyms CD3Z, T3Z, TCRZ

Function
Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR- mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:2470098, PubMed:7509083). CD3Z ITAMs phosphorylation creates multiple docking sites for the protein kinase ZAP70 leading to ZAP70 phosphorylation and its conversion into a catalytically active enzyme (PubMed:7509083). Plays an important role in intrathymic T-cell differentiation. Additionally, participates in the activity-dependent synapse formation of retinal ganglion cells (RGCs) in both the retina and dorsal lateral geniculate nucleus (dLGN) (By similarity).

Cellular Location
Cell membrane {ECO:0000250|UniProtKB:P24161}; Single-pass type I membrane protein

Tissue Location
CD3Z is expressed in normal lymphoid tissue and in peripheral blood mononuclear cells (PBMCs) (PubMed:11722641)

CD3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CD3 Antibody - Images




Figure 1: FCM analysis CD3 expressed in human peripheral blood cells.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999




Figure 1: FCM analysis CD3 expressed in human peripheral blood cells.

CD3 Antibody - References

1. Clevers,H.Annu.Rev.Immun.6:629-662,1988.