10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MMP3 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1661a

Specification

MMP3 Antibody - Product Information

Application	**E, WB, FC**
Primary Accession	P08254
Reactivity	**Human**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**IgG1**
Calculated MW	**54kDa KDa**

Description
Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. This gene encodes an enzyme which degrades fibronectin, laminin, collagens III, IV, IX, and X, and cartilage proteoglycans. The enzyme is thought to be involved in wound repair, progression of atherosclerosis, and tumor initiation. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3.

Immunogen
Purified recombinant fragment of human MMP3 expressed in E. Coli.

Formulation
Purified antibody in PBS with 0.05% sodium azide

MMP3 Antibody - Additional Information

Gene ID 4314

Other Names
Stromelysin-1, SL-1, 3.4.24.17, Matrix metalloproteinase-3, MMP-3, Transin-1, MMP3, STMY1

Dilution
E~~1/10000
WB~~1/500 - 1/2000
FC~~1/200 - 1/400

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
MMP3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MMP3 Antibody - Protein Information

Name MMP3

Synonyms STMY1

Function
Metalloproteinase with a rather broad substrate specificity that can degrade fibronectin, laminin, gelatins of type I, III, IV, and V; collagens III, IV, X, and IX, and cartilage proteoglycans. Activates different molecules including growth factors, plasminogen or other matrix metalloproteinases such as MMP9 (PubMed:11029580, PubMed:1371271). Once released into the extracellular matrix (ECM), the inactive pro-enzyme is activated by the plasmin cascade signaling pathway (PubMed:2383557). Acts also intracellularly (PubMed:22265821). For example, in dopaminergic neurons, gets activated by the serine protease HTRA2 upon stress and plays a pivotal role in DA neuronal degeneration by mediating microglial activation and alpha- synuclein/SNCA cleavage (PubMed:21330369). In addition, plays a role in immune response and possesses antiviral activity against various viruses such as vesicular stomatitis virus, influenza A virus (H1N1) and human herpes virus 1 (PubMed:35940311). Mechanistically, translocates from the cytoplasm into the cell nucleus upon virus infection to influence NF-kappa-B activities (PubMed:35940311).


Cellular Location
Secreted, extracellular space, extracellular matrix. Nucleus. Cytoplasm

MMP3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999






Figure 1: Western blot analysis using MMP3 mAb against human MMP3 (AA: 189-441) recombinant protein. (Expected MW is 54 kDa)




Figure 2: Flow cytometric analysis of NIH/3T3 cells using MMP3 mouse mAb (green) and negative control (red).

MMP3 Antibody - References

1. Int J Cancer. 2009 May 1;124(9):2172-8. 2. Cancer Epidemiol Biomarkers Prev. 2008 Dec;17(12):3551-7.