

#### MAPK8 Antibody

Purified Mouse Monoclonal Antibody Catalog # A01719a

# Specification

## MAPK8 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Calculated MW **Description**  E, WB, FC <u>P45983</u> Human, Mouse Mouse Monoclonal IgG1 48.3kDa KDa

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This kinase is activated by various cell stimuli, and targets specific transcription factors, and thus mediates immediate-early gene expression in response to cell stimuli. The activation of this kinase by tumor-necrosis factor alpha (TNF-alpha) is found to be required for TNF-alpha induced apoptosis. This kinase is also involved in UV radiation induced apoptosis, which is thought to be related to cytochrom c-mediated cell death pathway. Studies of the mouse counterpart of this gene suggested that this kinase play a key role in T cell proliferation, apoptosis and differentiation. Four alternatively spliced transcript variants encoding distinct isoforms have been reported.

Immunogen

Purified recombinant fragment of human MAPK8 expressed in E. Coli. <br />

**Formulation** Purified antibody in PBS with 0.05% sodium azide

### MAPK8 Antibody - Additional Information

Gene ID 5599

**Other Names** Mitogen-activated protein kinase 8, MAP kinase 8, MAPK 8, 2.7.11.24, JNK-46, Stress-activated protein kinase 1c, SAPK1c, Stress-activated protein kinase JNK1, c-Jun N-terminal kinase 1, MAPK8, JNK1, PRKM8, SAPK1, SAPK1C

**Dilution** E~~1/10000 WB~~1/500 - 1/2000 FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



### Precautions

MAPK8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### MAPK8 Antibody - Protein Information

#### Name MAPK8

### Function

Serine/threonine-protein kinase involved in various processes such as cell proliferation, differentiation, migration, transformation and programmed cell death. Extracellular stimuli such as pro- inflammatory cytokines or physical stress stimulate the stress- activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway (PubMed:<a

href="http://www.uniprot.org/citations/28943315" target=" blank">28943315</a>). In this cascade, two dual specificity kinases MAP2K4/MKK4 and MAP2K7/MKK7 phosphorylate and activate MAPK8/JNK1. In turn, MAPK8/JNK1 phosphorylates a number of transcription factors, primarily components of AP-1 such as JUN, JDP2 and ATF2 and thus regulates AP-1 transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/18307971" target=" blank">18307971</a>). Phosphorylates the replication licensing factor CDT1, inhibiting the interaction between CDT1 and the histone H4 acetylase HBO1 to replication origins (PubMed: <a href="http://www.uniprot.org/citations/21856198" target="\_blank">21856198</a>). Loss of this interaction abrogates the acetylation required for replication initiation (PubMed:<a href="http://www.uniprot.org/citations/21856198" target=" blank">21856198</a>). Promotes stressed cell apoptosis by phosphorylating key regulatory factors including p53/TP53 and Yesassociates protein YAP1 (PubMed: <a href="http://www.uniprot.org/citations/21364637" target=" blank">21364637</a>). In T-cells, MAPK8 and MAPK9 are required for polarized differentiation of T-helper cells into Th1 cells. Contributes to the survival of erythroid cells by phosphorylating the antagonist of cell death BAD upon EPO stimulation (PubMed:<a href="http://www.uniprot.org/citations/21095239" target="\_blank">21095239</a>). Mediates starvation-induced BCL2 phosphorylation, BCL2 dissociation from BECN1, and thus activation of autophagy (PubMed:<a href="http://www.uniprot.org/citations/18570871" target=" blank">18570871</a>). Phosphorylates STMN2 and hence regulates microtubule dynamics, controlling neurite elongation in cortical neurons (By similarity). In the developing brain, through its cytoplasmic activity on STMN2, negatively regulates the rate of exit from multipolar stage and of radial migration from the ventricular zone (By similarity). Phosphorylates several other substrates including heat shock factor protein 4 (HSF4), the deacetylase SIRT1, ELK1, or the E3 ligase ITCH (PubMed: <a href="http://www.uniprot.org/citations/20027304" target=" blank">20027304</a>, PubMed:<a href="http://www.uniprot.org/citations/16581800" target=" blank">16581800</a>, PubMed:<a href="http://www.uniprot.org/citations/17296730" target=" blank">17296730</a>). Phosphorylates the CLOCK-BMAL1 heterodimer and plays a role in the regulation of the circadian clock (PubMed:<a href="http://www.uniprot.org/citations/22441692" target=" blank">22441692</a>). Phosphorylates the heat shock transcription factor HSF1, suppressing HSF1-induced transcriptional activity (PubMed: <a href="http://www.uniprot.org/citations/10747973" target=" blank">10747973</a>). Phosphorylates POU5F1, which results in the inhibition of POU5F1's transcriptional activity and enhances its proteasomal degradation (By similarity). Phosphorylates JUND and this phosphorylation is inhibited in the presence of MEN1 (PubMed:<a href="http://www.uniprot.org/citations/22327296" target="\_blank">22327296</a>). In neurons, phosphorylates SYT4 which captures neuronal dense core vesicles at synapses (By similarity). Phosphorylates EIF4ENIF1/4-ET in response to oxidative stress, promoting P-body assembly (PubMed:<a href="http://www.uniprot.org/citations/22966201" target=" blank">22966201</a>).

Phosphorylates SIRT6 in response to oxidative stress, stimulating its mono-ADP-ribosyltransferase activity (PubMed:<a href="http://www.uniprot.org/citations/27568560"

target="\_blank">27568560</a>). Phosphorylates NLRP3, promoting assembly of the NLRP3 inflammasome (PubMed:<a href="http://www.uniprot.org/citations/28943315" target="\_blank">28943315</a>).



# **Cellular Location**

Cytoplasm. Nucleus. Synapse {ECO:0000250|UniProtKB:P49185}. Note=In the cortical neurons, predominantly cytoplasmic and associated with the Golgi apparatus and endosomal fraction. Increased neuronal activity increases phosphorylated form at synapses (By similarity). Colocalizes with POU5F1 in the nucleus. {ECO:0000250|UniProtKB:P49185, ECO:0000250|UniProtKB:Q91Y86}

## MAPK8 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>




Figure 1: Western blot analysis using MAPK8 mAb against human MAPK8 (AA: 227-380) recombinant protein. (Expected MW is 43.4 kDa)






Figure 2: Western blot analysis using MAPK8 mouse mAb against A431 (1), K562 (2), HeLa (3), NIH3T3 (4), PC-12 (5), and MCF-7 (6) cell lysate.




Figure 3: Flow cytometric analysis of HeLa cells using MAPK8 mouse mAb (green) and negative control (red).

### MAPK8 Antibody - References

1. Mol Biol Cell. 2011 Jan 1;22(1):117-27. 2. Cell Cycle. 2010 May 15;9(10):1951-9.