

SYCP3 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1855a

Specification

SYCP3 Antibody - Product Information

Application E, WB, IF, FC, IHC

Primary Accession
Reactivity
Human
Host
Clonality
Honoclonal
Isotype

Monoclonal

Calculated MW 27.7kDa KDa

Description

This gene encodes an essential structural component of the synaptonemal complex. This complex is involved in synapsis, recombination and segregation of meiotic chromosomes. Mutations in this gene are associated with azoospermia in males and susceptibility to pregnancy loss in females. Alternate splicing results in multiple transcript variants that encode the same protein.

Immunogen

Purified recombinant fragment of human SYCP3 (AA: 27-128) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

SYCP3 Antibody - Additional Information

Gene ID 50511

Other Names

Synaptonemal complex protein 3, SCP-3, SYCP3, SCP3

Dilution

E~~1/10000 WB~~1/500 - 1/2000 IF~~1/200 - 1/1000 FC~~1/200 - 1/400 IHC~~1/200 - 1/1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SYCP3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SYCP3 Antibody - Protein Information



Name SYCP3

Synonyms SCP3

Function

Component of the synaptonemal complexes (SCS), formed between homologous chromosomes during meiotic prophase. Required for centromere pairing during meiosis in male germ cells (By similarity). Required for normal meiosis during spermatogenesis and male fertility (PubMed:14643120). Plays a lesser role in female fertility. Required for efficient phosphorylation of HORMAD1 and HORMAD2 (By similarity).

Cellular Location


Nucleus {ECO:0000250|UniProtKB:Q60547}. Chromosome {ECO:0000250|UniProtKB:Q60547}. Chromosome, centromere {ECO:0000250|UniProtKB:Q60547}. Note=It is present in early unpaired cores, in the lateral domains of the synaptonemal complex and in the chromosome cores when they separate at diplotene. It is found axial to the metaphase I chromosomes and in association with pairs of sister centromeres. The centromere-associated protein becomes dissociated from the centromeres at anaphase II and is not found in mitotic metaphase centromeres. {ECO:0000250|UniProtKB:Q60547}

Tissue Location Testis-specific.

SYCP3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture








Figure 1: Western blot analysis using SYCP3 mAb against human SYCP3 recombinant protein. (Expected MW is 37.2 kDa)




Figure 2: Western blot analysis using SYCP3 mAb against HEK293 (1) and SYCP3 (AA: 27-128)-hlgGFc transfected HEK293 (2) cell lysate.




Figure 3: Immunofluorescence analysis of HepG2 cells using SYCP3 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.






Figure 4: Flow cytometric analysis of Jurkat cells using SYCP3 mouse mAb (green) and negative control (red).




Figure 5: Immunohistochemical analysis of paraffin-embedded endometrial cancer tissues using SYCP3 mouse mAb with DAB staining.





Figure 6: Immunohistochemical analysis of paraffin-embedded kidney tissues using SYCP3 mouse mAb with DAB staining.

SYCP3 Antibody - Background

This gene encodes an essential structural component of the synaptonemal complex. This complex is involved in synapsis, recombination and segregation of meiotic chromosomes. Mutations in this gene are associated with azoospermia in males and susceptibility to pregnancy loss in females. Alternate splicing results in multiple transcript variants that encode the same protein.

SYCP3 Antibody - References





1. Hum Pathol. 2013 Apr;44(4):472-9. 2. Cytogenet Genome Res. 2010;128(1-3):162-8.