

**Ring1 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO1861a****Specification****Ring1 Antibody - Product Information**

Application	E, WB, IF
Primary Accession	<a href="#">Q06587</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	42.4kDa KDa

**Description**

This gene belongs to the RING finger family, members of which encode proteins characterized by a RING domain, a zinc-binding motif related to the zinc finger domain. The gene product can bind DNA and can act as a transcriptional repressor. It is associated with the multimeric polycomb group protein complex. The gene product interacts with the polycomb group proteins BMI1, EDR1, and CBX4, and colocalizes with these proteins in large nuclear domains. It interacts with the CBX4 protein via its glycine-rich C-terminal domain. The gene maps to the HLA class II region, where it is contiguous with the RING finger genes FABGL and HKE4.

**Immunogen**

Purified recombinant fragment of human Ring1 (AA: 79-263) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**Ring1 Antibody - Additional Information**

**Gene ID** 6015

**Other Names**

E3 ubiquitin-protein ligase RING1, 6.3.2.-, Polycomb complex protein RING1, RING finger protein 1, Really interesting new gene 1 protein, RING1, RNF1

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IF~~1/200 - 1/1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Ring1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Ring1 Antibody - Protein Information

**Name** RING1 ([HGNC:10018](#))

### Function

Constitutes one of the E3 ubiquitin-protein ligases that mediate monoubiquitination of 'Lys-119' of histone H2A, thereby playing a central role in histone code and gene regulation. H2A 'Lys-119' ubiquitination gives a specific tag for epigenetic transcriptional repression and participates in X chromosome inactivation of female mammals. Essential component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones, rendering chromatin heritably changed in its expressibility. Compared to RNF2/RING2, it does not have the main E3 ubiquitin ligase activity on histone H2A, and it may rather act as a modulator of RNF2/RING2 activity.


### Cellular Location

Nucleus. Nucleus speckle

## Ring1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using Ring1 mAb against human Ring1 recombinant protein. (Expected MW is 44.6 kDa)




Figure 2: Western blot analysis using Ring1 mAb against HEK293 (1) and Ring1 (AA: 79-263)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Immunofluorescence analysis of HeLa cells using Ring1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 3: Immunofluorescence analysis of A431 cells using Ring1 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.

#### Ring1 Antibody - Background

The protein encoded by this gene plays a role in cell adhesion, and in cohesion of the endothelial monolayer at intercellular junctions in vascular tissue. Its expression may allow melanoma cells to interact with cellular elements of the vascular system, thereby enhancing hematogeneous tumor spread. Could be an adhesion molecule active in neural crest cells during ; embryonic development. Acts as surface receptor that triggers tyrosine phosphorylation of FYN and PTK2/FAK1, and a transient increase in the intracellular calcium concentration ;

#### Ring1 Antibody - References

1. Int J Dev Biol. 2009;53(2-3):355-70.
2. PLoS One. 2009 Dec 1;4(12):e8104.