

CDX2 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1885a**Specification****CDX2 Antibody - Product Information**

Application	E, WB, FC
Primary Accession	Q99626
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	33.5kDa KDa

Description

This gene is a member of the caudal-related homeobox transcription factor gene family. The encoded protein is a major regulator of intestine-specific genes involved in cell growth and differentiation. This protein also plays a role in early embryonic development of the intestinal tract. Aberrant expression of this gene is associated with intestinal inflammation and tumorigenesis.

Immunogen

Purified recombinant fragment of human CDX2 (AA: 176-303) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide

CDX2 Antibody - Additional Information

Gene ID 1045

Other Names

Homeobox protein CDX-2, CDX-3, Caudal-type homeobox protein 2, CDX2, CDX3

Dilution

E~~1/10000

WB~~1/500 - 1/2000

FC~~1/200 - 1/400

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CDX2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CDX2 Antibody - Protein Information

Name CDX2**Synonyms** CDX3**Function**

Transcription factor which regulates the transcription of multiple genes expressed in the intestinal epithelium (By similarity). Binds to the promoter of the intestinal sucrase-isomaltase SI and activates SI transcription (By similarity). Binds to the DNA sequence 5'-ATAAAACTTAT-3' in the promoter region of VDR and activates VDR transcription (By similarity). Binds to and activates transcription of LPH (By similarity). Activates transcription of CLDN2 and intestinal mucin MUC2 (By similarity). Binds to the 5'-AATTTTTTACAACACCT-3' DNA sequence in the promoter region of CA1 and activates CA1 transcription (By similarity). Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. Binds preferentially to methylated DNA (PubMed:28473536).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P43241}.


Tissue Location

Detected in small intestine, colon and pancreas.

CDX2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)






Figure 1: Western blot analysis using CDX2 mAb against human CDX2 (AA: 176-303) recombinant protein. (Expected MW is 40.1 kDa)




Figure 2: Western blot analysis using CDX2 mAb against HEK293 (1) and CDX2 (AA: 176-303)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Flow cytometric analysis of HeLa cells using CDX2 mouse mAb (green) and negative control (red).

CDX2 Antibody - Background

The protein encoded by this gene is a surface antigen that is preferentially expressed on monocytes/macrophages. It cooperates with other proteins to mediate the innate immune response to bacterial lipopolysaccharide. Alternative splicing results in multiple transcript variants encoding the same protein. ; ; ;

CDX2 Antibody - References

1. Tumour Biol. 2012 Dec;33(6):2185-8.
2. Cancer Biol Ther. 2012 Oct;13(12):1152-7.