

GABBR2 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO1902a**Specification****GABBR2 Antibody - Product Information**

Application	E, WB, IF, IHC
Primary Accession	O75899
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Calculated MW	105.8kDa KDa

Description

The multi-pass membrane protein encoded by this gene belongs to the G-protein coupled receptor 3 family and GABA-B receptor subfamily. The GABA-B receptors inhibit neuronal activity through G protein-coupled second-messenger systems, which regulate the release of neurotransmitters, and the activity of ion channels and adenylyl cyclase. This receptor subunit forms an active heterodimeric complex with GABA-B receptor subunit 1, neither of which is effective on its own. Allelic variants of this gene have been associated with nicotine dependence.

Immunogen

Purified recombinant fragment of human GABBR2 (AA: 319-483) expressed in E. Coli.

Formulation

Purified antibody in PBS with 0.05% sodium azide.

GABBR2 Antibody - Additional Information

Gene ID 9568

Other Names

Gamma-aminobutyric acid type B receptor subunit 2, GABA-B receptor 2, GABA-B-R2, GABA-BR2, GABABR2, Gb2, G-protein coupled receptor 51, HG20, GABBR2, GPR51, GPRC3B

Dilution

E~~1/10000
WB~~1/500 - 1/2000
IF~~1/200 - 1/1000
IHC~~1/200 - 1/1000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GABBR2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

GABBR2 Antibody - Protein Information

Name GABBR2

Synonyms GPR51, GPRC3B

Function

Component of a heterodimeric G-protein coupled receptor for GABA, formed by GABBR1 and GABBR2 (PubMed:9872316, PubMed:9872744, PubMed:15617512, PubMed:18165688, PubMed:22660477, PubMed:24305054). Within the heterodimeric GABA receptor, only GABBR1 seems to bind agonists, while GABBR2 mediates coupling to G proteins (PubMed:18165688). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase (PubMed:10075644, PubMed:10773016, PubMed:24305054). Signaling inhibits adenylate cyclase, stimulates phospholipase A2, activates potassium channels, inactivates voltage-dependent calcium-channels and modulates inositol phospholipid hydrolysis (PubMed:10075644, PubMed:9872744, PubMed:10906333, PubMed:10773016). Plays a critical role in the fine-tuning of inhibitory synaptic transmission (PubMed:9872744, PubMed:22660477). Pre-synaptic GABA receptor inhibits neurotransmitter release by down-regulating high-voltage activated calcium channels, whereas postsynaptic GABA receptor decreases neuronal excitability by activating a prominent inwardly rectifying potassium (Kir) conductance that underlies the late inhibitory postsynaptic potentials (PubMed:9872316, PubMed:10075644, PubMed:9872744, PubMed:22660477). Not only implicated in synaptic inhibition but also in hippocampal long-term potentiation, slow wave sleep, muscle relaxation and antinociception (Probable).

Cellular Location

Cell membrane; Multi-pass membrane protein. Postsynaptic cell membrane {ECO:0000250|UniProtKB:O88871}; Multi-pass membrane protein. Note=Coexpression of GABBR1 and GABBR2 is required for GABBR1 maturation and transport to the plasma membrane. In contrast, GABBR2 does not depend on GABBR1 for transport to the cell membrane

Tissue Location

Highly expressed in brain, especially in cerebral cortex, thalamus, hippocampus, frontal, occipital and temporal lobe, occipital pole and cerebellum, followed by corpus callosum, caudate nucleus, spinal cord, amygdala and medulla (PubMed:10087195, PubMed:10328880, PubMed:10727622, PubMed:9872744). Weakly expressed in heart, testis and skeletal muscle (PubMed:10087195, PubMed:10727622)

GABBR2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)





Figure 1: Western blot analysis using GABBR2 mAb against human GABBR2 (AA: 319-483) recombinant protein. (Expected MW is 44.9 kDa)




Figure 2: Western blot analysis using GABBR2 mAb against HEK293 (1) and GABBR2 (AA: 319-483)-hlgGfc transfected HEK293 (2) cell lysate.




Figure 3: Immunofluorescence analysis of HeLa cells using GABBR2 mouse mAb (green). Blue: DAPI fluorescent DNA dye. Secondary antibody from Fisher (Cat#: 35503)




Figure 4: Immunohistochemical analysis of paraffin-embedded cerebellum tissues using GABBR2 mouse mAb with DAB staining.

GABBR2 Antibody - Background

EIF2A is a 65-kD protein that catalyzes the formation of puromycin-sensitive 80S preinitiation

complexes ; ;

GABBR2 Antibody - References

1. J Biol Chem. 2007 Feb 9;282(6):4162-71.
2. Genomics. 1999 Mar 15;56(3):288-95.