

TUBE1 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO1948a

Specification

TUBE1 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Calculated MW Description E, WB, FC <u>09UJT0</u> Human Mouse Monoclonal IgG1 53kDa KDa

This gene encodes a member of the tubulin superfamily. This protein localizes to the centriolar sub-distal appendages that are associated with the older of the two centrioles after centrosome duplication. This protein plays a central role in organization of the microtubules during centriole duplication. A pseudogene of this gene is found on chromosome 5.

Immunogen Purified recombinant fragment of human TUBE1 (AA: 314-472) expressed in E. Coli.

Formulation Purified antibody in PBS with 0.05% sodium azide.

TUBE1 Antibody - Additional Information

Gene ID 51175

Other Names Tubulin epsilon chain, Epsilon-tubulin, TUBE1, TUBE

Dilution E~~1/10000 WB~~1/500 - 1/2000 FC~~1/200 - 1/400

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions TUBE1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TUBE1 Antibody - Protein Information

Name TUBE1



Synonyms TUBE

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Note=Associated with pericentriolar material.

TUBE1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>




Figure 1: Western blot analysis using TUBE1 mAb against human TUBE1 (AA: 314-472) recombinant protein. (Expected MW is 44.3 kDa)






Figure 2: Western blot analysis using TUBE1 mAb against HEK293 (1) and TUBE1 (AA: 314-472)-hlgGFc transfected HEK293 (2) cell lysate.




Figure 3: Flow cytometric analysis of Hela cells using TUBE1 mouse mAb (green) and negative control (red).

TUBE1 Antibody - Background

The protein encoded by this gene belongs to the cyclic nucleotide phosphodiesterase (PDE) family, and PDE1 subfamily. Members of the PDE1 family are calmodulin-dependent PDEs that are stimulated by a calcium-calmodulin complex. This PDE has dual-specificity for the second messengers, cAMP and cGMP, with a preference for cGMP as a substrate. cAMP and cGMP function as key regulators of many important physiological processes. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. ; ;

TUBE1 Antibody - References

1. Nat Cell Biol. 2003 Jan;5(1):71-6.2. Nat Cell Biol. 2000 Jan;2(1):30-5.