

**WTAP Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO2050a****Specification****WTAP Antibody - Product Information**

Application	E, WB, IF, FC, IHC
Primary Accession	<a href="#">Q15007</a>
Reactivity	Human, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	44.2kDa KDa

**Description**

The Wilms tumor suppressor gene WT1 appears to play a role in both transcriptional and posttranscriptional regulation of certain cellular genes. This gene encodes a WT1-associating protein, which is a ubiquitously expressed nuclear protein. Like WT1 protein, this protein is localized throughout the nucleoplasm as well as in speckles and partially colocalizes with splicing factors. Alternative splicing of this gene results in several transcript variants encoding three different isoforms.

**Immunogen**

Purified recombinant fragment of human WTAP (AA: 91-201) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**WTAP Antibody - Additional Information**

**Gene ID** 9589

**Other Names**

Pre-mRNA-splicing regulator WTAP, Female-lethal(2)D homolog, hFL(2)D, WT1-associated protein, Wilms tumor 1-associating protein, WTAP, KIAA0105

**Dilution**

E~~1/10000  
WB~~1/500 - 1/2000  
IF~~1/200 - 1/1000  
FC~~1/200 - 1/400  
IHC~~1/200 - 1/1000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

WTAP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## WTAP Antibody - Protein Information

**Name** WTAP {ECO:0000303|PubMed:11001926, ECO:0000312|HGNC:HGNC:16846}

### Function

Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing (PubMed:<a href="http://www.uniprot.org/citations/29507755" target="\_blank">29507755</a>). Required for accumulation of METTL3 and METTL14 to nuclear speckle (PubMed:<a href="http://www.uniprot.org/citations/24316715" target="\_blank">24316715</a>, PubMed:<a href="http://www.uniprot.org/citations/24407421" target="\_blank">24407421</a>, PubMed:<a href="http://www.uniprot.org/citations/24981863" target="\_blank">24981863</a>). Acts as a mRNA splicing regulator (PubMed:<a href="http://www.uniprot.org/citations/12444081" target="\_blank">12444081</a>). Regulates G2/M cell-cycle transition by binding to the 3' UTR of CCNA2, which enhances its stability (PubMed:<a href="http://www.uniprot.org/citations/17088532" target="\_blank">17088532</a>). Impairs WT1 DNA-binding ability and inhibits expression of WT1 target genes (PubMed:<a href="http://www.uniprot.org/citations/17095724" target="\_blank">17095724</a>).

### Cellular Location

Nucleus speckle. Nucleus, nucleoplasm. Cytoplasm {ECO:0000250|UniProtKB:Q9ER69}. Note=Mainly nuclear with some fraction located in the cytoplasm. ZC3H13 is required to anchor component of the MACOM subcomplex, such as VIRMA, in the nucleus {ECO:0000250|UniProtKB:Q9ER69}

### Tissue Location

Ubiquitously expressed.

## WTAP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)