

#### Notch3 Antibody

Purified Mouse Monoclonal Antibody Catalog # AO2126a

# Specification

## Notch3 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Isotype Calculated MW **Description**  E, WB <u>O9UM47</u> Human Mouse Monoclonal IgG2b 243.6kDa KDa

This gene encodes the third discovered human homologue of the Drosophilia melanogaster type I membrane protein notch. In Drosophilia, notch interaction with its cell-bound ligands (delta, serrate) establishes an intercellular signalling pathway that plays a key role in neural development. Homologues of the notch-ligands have also been identified in human, but precise interactions between these ligands and the human notch homologues remains to be determined. Mutations in NOTCH3 have been identified as the underlying cause of cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL).

Immunogen Purified recombinant fragment of human Notch3 (AA: 47-156) expressed in E. Coli.

**Formulation** Purified antibody in PBS with 0.05% sodium azide

### **Notch3 Antibody - Additional Information**

Gene ID 4854

**Other Names** Neurogenic locus notch homolog protein 3, Notch 3, Notch 3 extracellular truncation, Notch 3 intracellular domain, NOTCH3

**Dilution** E~~1/10000 WB~~1/500 - 1/2000

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Notch3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **Notch3 Antibody - Protein Information**



## Name NOTCH3

### Function

Functions as a receptor for membrane-bound ligands Jagged1, Jagged2 and Delta1 to regulate cell-fate determination (PubMed:<a href="http://www.uniprot.org/citations/15350543" target="\_blank">15350543</a>). Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBPJ/RBPSUH and activates genes of the enhancer of split locus. Affects the implementation of differentiation, proliferation and apoptotic programs (By similarity).

**Cellular Location** 

Cell membrane; Single-pass type I membrane protein

**Tissue Location** 

Ubiquitously expressed in fetal and adult tissues.

# Notch3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>