

**CBX2 Antibody**  
**Purified Mouse Monoclonal Antibody**  
**Catalog # AO2138a****Specification****CBX2 Antibody - Product Information**

Application	E, WB
Primary Accession	<a href="#">Q14781</a>
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	56kDa KDa

**Description**

This gene encodes a component of the polycomb multiprotein complex, which is required to maintain the transcriptionally repressive state of many genes throughout development via chromatin remodeling and modification of histones. Disruption of this gene in mice results in male-to-female gonadal sex reversal. Mutations in this gene are also associated with gonadal dysgenesis in humans. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene.

**Immunogen**

Purified recombinant fragment of human CBX2 (AA: 402-525) expressed in E. Coli.

**Formulation**

Purified antibody in PBS with 0.05% sodium azide

**CBX2 Antibody - Additional Information**

**Gene ID** 84733

**Other Names**

Chromobox protein homolog 2, CBX2

**Dilution**

E~~1/10000

WB~~1/500 - 1/2000

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CBX2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CBX2 Antibody - Protein Information**

**Name** CBX2**Function**

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development (PubMed:<a href="http://www.uniprot.org/citations/21282530" target="\_blank">21282530</a>). PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:<a href="http://www.uniprot.org/citations/21282530" target="\_blank">21282530</a>). Binds to histone H3 trimethylated at 'Lys-9' (H3K9me3) or at 'Lys-27' (H3K27me3) (By similarity). Plays a role in the lineage differentiation of the germ layers in embryonic development (By similarity). Involved in sexual development, acting as activator of NR5A1 expression (PubMed:<a href="http://www.uniprot.org/citations/19361780" target="\_blank">19361780</a>).

**Cellular Location**

Nucleus. Chromosome Note=Localized in distinct foci on chromatin and in chromocenters Localizes to the inactive X chromosome. Seems to be recruited to H3K27me3, H3K9ac and H3K3me2 sites on chromatin

**CBX2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)