10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SLC2A4 Antibody
Purified Mouse Monoclonal Antibody
Catalog # AO2244a

Specification

SLC2A4 Antibody - Product Information

Application	**E, WB, IF, FC, IHC**
Primary Accession	P14672
Reactivity	**Human, Mouse**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**IgG2b**
Calculated MW	**54.8kDa KDa**

Description
This gene is a member of the solute carrier family 2 (facilitated glucose transporter) family and encodes a protein that functions as an insulin-regulated facilitative glucose transporter. In the absence of insulin, this integral membrane protein is sequestered within the cells of muscle and adipose tissue. Within minutes of insulin stimulation, the protein moves to the cell surface and begins to transport glucose across the cell membrane. Mutations in this gene have been associated with noninsulin-dependent diabetes mellitus (NIDDM).

Immunogen
Purified recombinant fragment of human SLC2A4 (AA: 224-353) expressed in E. Coli.

Formulation
Ascitic fluid containing 0.03% sodium azide.

SLC2A4 Antibody - Additional Information

Gene ID 6517

Other Names
Solute carrier family 2, facilitated glucose transporter member 4, Glucose transporter type 4, insulin-responsive, GLUT-4, SLC2A4, GLUT4

Dilution
E~~1/10000
WB~~1/500 - 1/2000
IF~~1/200 - 1/1000
FC~~1/200 - 1/400
IHC~~1/200 - 1/1000

Storage
Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
SLC2A4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

SLC2A4 Antibody - Protein Information

Name SLC2A4 (HGNC:11009)

Function
Insulin-regulated facilitative glucose transporter, which plays a key role in removal of glucose from circulation. Response to insulin is regulated by its intracellular localization: in the absence of insulin, it is efficiently retained intracellularly within storage compartments in muscle and fat cells. Upon insulin stimulation, translocates from these compartments to the cell surface where it transports glucose from the extracellular milieu into the cell.


Cellular Location
Cell membrane {ECO:0000250|UniProtKB:P14142}; Multi-pass membrane protein {ECO:0000250|UniProtKB:P14142} Endomembrane system; Multi-pass membrane protein. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:P14142}. Note=Localizes primarily to the perinuclear region, undergoing continued recycling to the plasma membrane where it is rapidly reinternalized (PubMed:8300557). The dileucine internalization motif is critical for intracellular sequestration (PubMed:8300557). Insulin stimulation induces translocation to the cell membrane (By similarity) {ECO:0000250|UniProtKB:P14142, ECO:0000269|PubMed:8300557}

Tissue Location
Skeletal and cardiac muscles; brown and white fat.

SLC2A4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999




Figure 1: Western blot analysis using SLC2A4 mAb against human SLC2A4 recombinant protein. (Expected MW is 39.9 kDa)




Figure 2: Western blot analysis using SLC2A4 mAb against HEK293 (1) and SLC2A4 (AA: 224-353)-hIgGFc transfected HEK293 (2) cell lysate.




Figure 3: Western blot analysis using SLC2A4 mouse mAb against NIH3T3 (1), 3T3L1 (2), MCF-7 (4) cell lysate and Mouse heart (3) tissue lysate.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999




Figure 4: Immunofluorescence analysis of HeLa cells using SLC2A4 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 5: Immunofluorescence analysis of HepG2 cells using SLC2A4 mouse mAb (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.




Figure 6: Flow cytometric analysis of HeLa cells using SLC2A4 mouse mAb (green) and negative control (purple).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999




Figure 7: Immunohistochemical analysis of paraffin-embedded bladder cancer tissues using SLC2A4 mouse mAb with DAB staining.




Figure 8: Immunohistochemical analysis of paraffin-embedded cardiac muscle tissues using SLC2A4 mouse mAb with DAB staining.

SLC2A4 Antibody - References

1.J Biol Chem. 2011 May 13;286(19):16541-5. 2.PLoS One. 2010 Dec 20;5(12):e15560.