10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CO039 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10000B

Specification

CO039 Antibody (C-term) - Product Information

Application	**WB, FC,E**
Primary Accession	Q6ZRI6
Other Accession	NP_056307.2
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**110673**
Antigen Region	**938-967**

CO039 Antibody (C-term) - Additional Information

Gene ID 56905

Other Names
Uncharacterized protein C15orf39, C15orf39

Target/Specificity
This CO039 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 938-967 amino acids from the C-terminal region of human CO039.

Dilution
WB~~1:1000
FC~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
CO039 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CO039 Antibody (C-term) - Protein Information

Name C15orf39


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

CO039 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CO039 Antibody (C-term) - Images



CO039 Antibody (C-term) (Cat. #AP10000b) western blot analysis in A549 cell line lysates (35ug/lane).This demonstrates the CO039 antibody detected the CO039 protein (arrow).



CO039 Antibody (C-term) (Cat. #AP10000b) flow cytometric analysis of A549 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.