

### **POTEB Antibody (C-term)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10032b

#### Specification

## **POTEB Antibody (C-term) - Product Information**

Application Primary Accession Other Accession

Reactivity Host Clonality Isotype Antigen Region WB, IHC-P, FC,E <u>O6S5H4</u> <u>POCG39</u>, <u>POCG38</u>, <u>A5A3E0</u>, <u>O86YR6</u>, <u>B2RU33</u>, <u>H3BUK9</u>, <u>NP\_997238.2</u> Human Rabbit Polyclonal Rabbit IgG 381-410

### **POTEB** Antibody (C-term) - Additional Information

**Other Names** 

POTE ankyrin domain family member B, ANKRD26-like family B member 1, Prostate, ovary, testis-expressed protein on chromosome 15, POTE-15, POTEB, A26B1, POTE15

Target/Specificity

This POTEB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 381-410 amino acids from the C-terminal region of human POTEB.

**Dilution** WB~~1:2000 IHC-P~~1:50~100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** 

POTEB Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

### **POTEB Antibody (C-term) - Protein Information**


### **POTEB Antibody (C-term) - Protocols**



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# POTEB Antibody (C-term) - Images




All lanes : Anti-POTEB Antibody (C-term) at 1:2000 dilution Lane 1: Human ovary tissue lysate Lane 2: Human liver tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 66 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



POTEB antibody (C-term) (Cat. #AP10032b) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the POTEB antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.





POTEB Antibody (C-term) (Cat. #AP10032b) flow cytometric analysis of Jurkat cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.