10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

MICALL1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10498a

Specification

MICALL1 Antibody (C-term) - Product Information

Application	**WB, IHC-P,E**
Primary Accession	Q8N3F8
Other Accession	NP_203744.1
Reactivity	**Human**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**93441**
Antigen Region	**752-781**

MICALL1 Antibody (C-term) - Additional Information

Gene ID 85377

Other Names
MICAL-like protein 1, Molecule interacting with Rab13, MIRab13, MICALL1, KIAA1668, MIRAB13

Target/Specificity
This MICALL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 752-781 amino acids from the C-terminal region of human MICALL1.

Dilution
WB~~1:1000
IHC-P~~1:50~100

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
MICALL1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MICALL1 Antibody (C-term) - Protein Information

Name MICALL1

Synonyms KIAA1668, MIRAB13

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Function Probable lipid-binding protein with higher affinity for phosphatidic acid, a lipid enriched in recycling endosome membranes. On endosome membranes, may act as a downstream effector of Rab proteins recruiting cytosolic proteins to regulate membrane tubulation. May be involved in a late step of receptor-mediated endocytosis regulating for instance endocytosed-EGF receptor trafficking. Alternatively, may regulate slow endocytic recycling of endocytosed proteins back to the plasma membrane. May indirectly play a role in neurite outgrowth.


Cellular Location
Recycling endosome membrane; Peripheral membrane protein. Late endosome membrane. Note=Localization to late endosomes is actin-dependent. Association to tubular recycling endosomes is regulated by RAB35 and ARF6

MICALL1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MICALL1 Antibody (C-term) - Images



MICALL1 Antibody (C-term) (Cat. #AP10498a) western blot analysis in U251 cell line lysates (35ug/lane).This demonstrates the MICALL1 antibody detected the LACRT protein (arrow).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



MICALL1 Antibody (C-term) (Cat. #AP10498a) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MICALL1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

MICALL1 Antibody (C-term) - Background

MICALL1 may be a cytoskeletal regulator.

MICALL1 Antibody (C-term) - References

Sharma, M., et al. Mol. Biol. Cell 20(24):5181-5194(2009)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Jin, J., et al. Curr. Biol. 14(16):1436-1450(2004)
Beausoleil, S.A., et al. Proc. Natl. Acad. Sci. U.S.A. 101(33):12130-12135(2004)