

KIAA1609 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10514b**Specification**

KIAA1609 Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q6P9B6
Other Accession	NP_065998.3
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	426-456

KIAA1609 Antibody (C-term) - Additional Information**Gene ID** 57707**Other Names**

TLD domain-containing protein 1, TBC/LysM-associated domain-containing protein 1, TLDC1, KIAA1609

Target/Specificity

This KIAA1609 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 426-456 amino acids from the C-terminal region of human K1609.

DilutionWB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

KIAA1609 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

KIAA1609 Antibody (C-term) - Protein Information**Name** MEAK7 ([HGNC:29325](#))

Synonyms KIAA1609, TLDC1

Function Activates an alternative mTOR signaling through RPS6KB2 activation and EIF4EBP1 repression to regulate cell proliferation and migration (PubMed:[29750193](#)). Recruits MTOR at the lysosome, essential for MTOR signaling at the lysosome (PubMed:[29750193](#)).

Cellular Location


Membrane. Cytoplasm. Lysosome

KIAA1609 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


KIAA1609 Antibody (C-term) - Images




Anti-KIAA1609 Antibody (C-term) at 1:8000 dilution + DU145 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



KIAA1609 Antibody (C-term) (Cat. #AP10514b) western blot analysis in mouse NIH-3T3 cell line lysates (35ug/lane). This demonstrates the KIAA1609 antibody detected the K1609 protein (arrow).



KIAA1609 Antibody (C-term) (Cat. #AP10514b) immunohistochemistry analysis in formalin fixed and paraffin embedded human uterus tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the KIAA1609 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



KIAA1609 Antibody (C-term) (Cat. #AP10514b) flow cytometric analysis of NIH-3T3 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit

secondary antibodies were used for the analysis.

KIAA1609 Antibody (C-term) - References

Rothenberg, M.E., et al. Nat. Genet. 42(4):289-291(2010)

Venkatesan, K., et al. Nat. Methods 6(1):83-90(2009)

Wang, A.G., et al. Biochem. Biophys. Res. Commun. 345(3):1022-1032(2006)