

## **5HT3E Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10528c

### **Specification**

## **5HT3E Antibody (Center) - Product Information**

WB, FC, E Application **Primary Accession** A5X5Y0 Other Accession NP 872395.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 51438 Antigen Region 126-154

### **5HT3E Antibody (Center) - Additional Information**

#### Gene ID 285242

#### **Other Names**

5-hydroxytryptamine receptor 3E, 5-HT3-E, 5-HT3E, Serotonin receptor 3E, HTR3E

#### Target/Specificity

This 5HT3E antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 126-154 amino acids from the Central region of human 5HT3E.

# **Dilution**

WB~~1:1000 FC~~1:10~50

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

5HT3E Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### **5HT3E Antibody (Center) - Protein Information**

## Name HTR3E (HGNC:24005)

Function Forms serotonin (5-hydroxytryptamine/5-HT3)-activated cation- selective channel





Tel: 858.875.1900 Fax: 858.875.1999

complexes, which when activated cause fast, depolarizing responses in neurons.

#### **Cellular Location**

Postsynaptic cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=Presumably retained within the endoplasmic reticulum unless complexed with HTR3A.

#### **Tissue Location**


Expressed in adult colon and intestine.

## **5HT3E Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### 5HT3E Antibody (Center) - Images



5HT3E Antibody (Center) (Cat. #AP10528c) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the 5HT3E antibody detected the 5HT3E protein (arrow).





5HT3E Antibody (Center) (Cat. #AP10528c) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# 5HT3E Antibody (Center) - Background

The product of this gene belongs to the ligand-gated ion channel receptor superfamily. HTR3E encodes a subunit E of the type 3 receptor for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. This receptor causes fast, depolarizing responses in neurons after activation. Genes encoding subunits C, D and E form a cluster on chromosome 3. An alternative splice variant has been described but its full length sequence has not been determined.

### 5HT3E Antibody (Center) - References

Walstab, J., et al. J. Biol. Chem. 285(35):26956-26965(2010)
Hammer, C., et al. Pharmacogenomics 11(7):943-950(2010)
Lennertz, L., et al. Eur Neuropsychopharmacol 20(6):414-420(2010)
Goecke, T.W., et al. Acta Obstet Gynecol Scand 89(1):7-14(2010)
Schuhmacher, A., et al. Pharmacogenet. Genomics 19(11):843-851(2009)