

ZFYVE28 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP10577b

Specification

ZFYVE28 Antibody (C-term) - Product Information

Application IHC-P,E Primary Accession O9HCC9

Other Accession <u>NP_001166130.1</u>, <u>NP_066023.2</u>,

NP 001166131.1, NP 001166127.1,

NP 001166129.1

Reactivity
Human
Host
Clonality
Polyclonal
Isotype
Antigen Region
Rabbit IgG
604-633

ZFYVE28 Antibody (C-term) - Additional Information

Gene ID 57732

Other Names

Lateral signaling target protein 2 homolog, hLst2, Zinc finger FYVE domain-containing protein 28, ZFYVE28, KIAA1643, LST2

Target/Specificity

This ZFYVE28 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 604-633 amino acids from the C-terminal region of human ZFYVE28.

Dilution

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZFYVE28 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZFYVE28 Antibody (C-term) - Protein Information

Name ZFYVE28



Synonyms KIAA1643, LST2

Function Negative regulator of epidermal growth factor receptor (EGFR) signaling. Acts by promoting EGFR degradation in endosomes when not monoubiquitinated.

Cellular Location

Cytoplasm, cytosol. Early endosome membrane. Note=Localizes to early endosome membrane in absence of Lys-87 monoubiquitination Localizes to cytosol when monoubiquitinated

ZFYVE28 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ZFYVE28 Antibody (C-term) - Images



ZFYVE28 antibody (C-term) (Cat. #AP10577b) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ZFYVE28 antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

ZFYVE28 Antibody (C-term) - Background

Negative regulator of epidermal growth factor receptor (EGFR) signaling. Acts by promoting EGFR degradation in endosomes when not monoubiquitinated.