

FBXW8 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10719c**Specification**

FBXW8 Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q8N3Y1
Other Accession	NP_036306.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	270-299

FBXW8 Antibody (Center) - Additional Information**Gene ID** 26259**Other Names**

F-box/WD repeat-containing protein 8, F-box and WD-40 domain-containing protein 8, F-box only protein 29, FBXW8, FBW6, FBW8, FBX29, FBXO29, FBXW6

Target/Specificity

This FBXW8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 270-299 amino acids from the Central region of human FBXW8.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FBXW8 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

FBXW8 Antibody (Center) - Protein Information**Name** FBXW8

Synonyms FBW6, FBW8, FBX29, FBXO29, FBXW6

Function Substrate-recognition component of a Cul7-RING ubiquitin- protein ligase complex, which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. The Cul7- RING(FBXW8) complex mediates ubiquitination and consequent degradation of GORASP1, acting as a component of the ubiquitin ligase pathway that regulates Golgi morphogenesis and dendrite patterning in brain (PubMed:[21572988](#)). Mediates ubiquitination and degradation of IRS1 in a mTOR-dependent manner: the Cul7-RING(FBXW8) complex recognizes and binds IRS1 previously phosphorylated by S6 kinase (RPS6KB1 or RPS6KB2) (PubMed:[18498745](#)). The Cul7-RING(FBXW8) complex also mediates ubiquitination of MAP4K1/HPK1: recognizes and binds autophosphorylated MAP4K1/HPK1, leading to its degradation, thereby affecting cell proliferation and differentiation (PubMed:[24362026](#)). Associated component of the 3M complex, suggesting that it mediates some of 3M complex functions (PubMed:[24793695](#)).

Cellular Location


Cytoplasm, perinuclear region. Golgi apparatus. Note=Colocalizes with CUL7 at the Golgi apparatus in neurons. {ECO:0000250|UniProtKB:P0DL28}

FBXW8 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


FBXW8 Antibody (Center) - Images




FBXW8 Antibody (Center) (Cat. #AP10719c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the FBXW8 antibody detected the FBXW8 protein (arrow).



FBXW8 Antibody (Center) (Cat. #AP10719c) western blot analysis in mouse bladder tissue lysates (35ug/lane). This demonstrates the FBXW8 antibody detected the FBXW8 protein (arrow).



FBXW8 Antibody (Center) (Cat. #AP10719c) immunohistochemistry analysis in formalin fixed and paraffin embedded human lung carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the FBXW8 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



FBXW8 Antibody (Center) (Cat. #AP10719c) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

FBXW8 Antibody (Center) - Background

This gene encodes a member of the F-box protein family, members of which are characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into three classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene contains a WD-40 domain, in addition to an F-box motif, so it belongs to the Fbw class. Alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene.

FBXW8 Antibody (Center) - References

Tsutsumi, T., et al. Mol. Cell. Biol. 28(2):743-751(2008)
Koch, H.B., et al. Cell Cycle 6(2):205-217(2007)
Okabe, H., et al. PLoS ONE 1, E128 (2006) :
Watanabe, N., et al. Proc. Natl. Acad. Sci. U.S.A. 101(13):4419-4424(2004)
Dias, D.C., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16601-16606(2002)

FBXW8 Antibody (Center) - Citations

- [The CUL7/F-box and WD repeat domain containing 8 \(CUL7/Fbxw8\) ubiquitin ligase promotes degradation of hematopoietic progenitor kinase 1.](#)