10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HIST1H2BJ Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP10727c

Specification

HIST1H2BJ Antibody (Center) - Product Information

Application	**WB, IHC-P, FC,E**
Primary Accession	P06899
Other Accession	Q9PSW9, P0C1H5, P0C1H4, Q6PC60, Q5QNW6, Q16778, Q6DRA6, Q6DN03, Q64525, Q5BJA5, P0C1H3, P62808, P23527, Q99877, Q32L48, P10854, Q99879, Q2PFX4, O60814, Q2M2T1, Q64478, Q93079, P58876, Q6ZWY9, P62807, P33778, Q00729, P06900, P02281, NP_066402.2
Reactivity	**Human**
Predicted	**Xenopus, Rat, Mouse, Bovine, Monkey, Chicken, Zebrafish**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Antigen Region	**57-86**

HIST1H2BJ Antibody (Center) - Additional Information

Gene ID 8970

Other Names
Histone H2B type 1-J, Histone H2B1, Histone H2Br, H2B/r, HIST1H2BJ, H2BFR

Target/Specificity
This HIST1H2B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 57-86 amino acids from the Central region of human HIST1H2BJ.

Dilution
WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
HIST1H2BJ Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HIST1H2BJ Antibody (Center) - Protein Information

Name H2BC11 (HGNC:4761)

Function Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


Cellular Location
Nucleus. Chromosome.

HIST1H2BJ Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HIST1H2BJ Antibody (Center) - Images




HIST1H2B Antibody (Center) (Cat. #AP10727c) western blot analysis in CEM,HL-60,NCI-H460 cell line lysates (35ug/lane).This demonstrates the HIST1H2B antibody detected the HIST1H2B protein (arrow).


10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western blot analysis of lysates from ZF4 cell line, Zebrafish, zebra fish muscle tissue lysate(from left to right), using HIST1H2BJ Antibody (Center)(Cat. #AP10727c). AP10727c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



HIST1H2B antibody (Center) (Cat. #AP10727c) immunohistochemistry analysis in formalin fixed and paraffin embedded human hepatocarcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the HIST1H2B antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

HIST1H2B Antibody (Center) (Cat. #AP10727c) flow cytometric analysis of CEM cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HIST1H2BJ Antibody (Center) - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a member of the histone H2B family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq].

HIST1H2BJ Antibody (Center) - References

Shi, J., et al. Nature 460(7256):753-757(2009)
Benyamin, B., et al. Am. J. Hum. Genet. 84(1):60-65(2009)
Kim, S.C., et al. Mol. Cell 23(4):607-618(2006)
Beck, H.C., et al. Mol. Cell Proteomics 5(7):1314-1325(2006)
Pavri, R., et al. Cell 125(4):703-717(2006)

HIST1H2BJ Antibody (Center) - Citations

- Distinct Motion of GFP-Tagged Histone Expressing Cells Under AC Electrokinetics in Electrode-Multilayered Microfluidic Device.